

Free products of finite dimensional and other von Neumann algebras with respect to non-tracial states.

Kenneth J. Dykema

Department of Mathematics
University of California
Berkeley CA 94720-3840

present address:

Department of Mathematics and Computer Science
Odense Universitet, Campusvej 55
DK-5230 Odense M, Denmark
e-mail: dykema@imada.ou.dk

Abstract. The von Neumann algebra free product of arbitrary finite dimensional von Neumann algebras with respect to arbitrary faithful states, at least one of which is not a trace, is found to be a type III factor possibly direct sum a finite dimensional algebra. The free product state on the type III factor is what we call an extremal almost periodic state, and has centralizer isomorphic to $L(F_\infty)$. This allows further classification the type III factor and provides another construction of full type III₁ factors having arbitrary Sd invariant of Connes. The free products considered in this paper are not limited to free products of finite dimensional algebras, but can be of a quite general form.

Introduction.

In the context of Voiculescu's theory of freeness, (see Voiculescu, Dykema and Nica [1992]), the free product of operator algebras is a basic operation, which is to freeness what the tensor product is to independence. Free products of several sorts of von Neumann algebras have been investigated, including free products of finite dimensional and hyperfinite von Neumann algebras with respect to traces, (Voiculescu [1990], Dykema [1993(a)], Dykema [1994(a)], Dykema [1993(b)]), certain amalgamated free products with respect to traces, (Popa [1993], Rădulescu [1994], Dykema [1995(a)]) and free products of various von Neumann algebras with respect to non-tracial states, (Rădulescu [preprint], Barnett [1995], Dykema [1994(b)]). In Rădulescu [preprint], the free product of the diffuse abelian von Neumann algebra and $M_2(\mathbf{C})$ with respect to the state $\text{Tr} \left(\begin{pmatrix} \frac{1}{1+\lambda} & 0 \\ 0 & \frac{\lambda}{1+\lambda} \end{pmatrix} \cdot \right)$ for $0 < \lambda < 1$ was shown to be a type III_λ factor with core $L(F_\infty) \otimes B(\mathcal{H})$. In Barnett [1995] and Dykema [1994(b)], some general results about free products of von Neumann algebras with respect to non-tracial states were proved, but several natural questions were left unanswered. For example, in the free product

$$(\mathcal{M}, \phi) = (M_2(\mathbf{C}), \text{Tr} \left(\begin{pmatrix} \frac{1}{1+\lambda} & 0 \\ 0 & \frac{\lambda}{1+\lambda} \end{pmatrix} \cdot \right)) * (M_2(\mathbf{C}), \text{Tr} \left(\begin{pmatrix} \frac{1}{1+\mu} & 0 \\ 0 & \frac{\mu}{1+\mu} \end{pmatrix} \cdot \right))$$

Supported by a National Science Foundation Postdoctoral Fellowship and by a Fields Institute Fellowship.

for $0 < \lambda, \mu < 1$, $\mathcal{M}$ was by Dykema [1994(b)] known to be a factor only for certain values of λ and μ , and for other values this remained unknown. One of the consequences of our main result is that such a free product is always a factor, and the centralizer of the free product state ϕ is isomorphic to $L(F_\infty)$.

In this paper, we investigate the free product of finite dimensional von Neumann algebras (and others) with respect to arbitrary faithful states that are not traces. The main result can be summarized as follows:

Theorem 1 *Let*

$$(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2) \quad (1)$$

be the von Neumann algebra free product of finite dimensional algebras with respect to faithful states, at least one of which is non-tracial. Then

$$\mathcal{M} = \mathcal{M}_0 \oplus D \quad \text{or} \quad \mathcal{M} = \mathcal{M}_0, \quad (2)$$

where D is a finite dimensional algebra and where $\mathcal{M}_0$ is a type III factor. The type I part, D , can be found from knowledge of ϕ_1 and ϕ_2 , as described below. The restriction, $\phi_0 \stackrel{\text{def}}{=} \phi|_{\mathcal{M}_0}$ to the type III part, is an almost periodic state (or functional) whose centralizer is isomorphic to the II_1 -factor $L(F_\infty)$. The point spectrum of the modular operator of ϕ_0 , Δ_{ϕ_0} , is equal to the subgroup of $\mathbf{R}_+^*$ generated by the union of the point spectra of Δ_{ϕ_1} and of Δ_{ϕ_2} . Thus, in Connes' classification of $\mathcal{M}_0$ as type III_λ , one can find λ and always $0 < \lambda \leq 1$.

Note that in (1) when both ϕ_1 and ϕ_2 are traces, $\mathcal{M}$ was found in Dykema [1993(b)]. In that case, $\mathcal{M}$ was of a similar form, except that the non-(type I) part, $\mathcal{M}_0$, was then a II_1 factor related to free groups, (see Rădulescu [1994] and Dykema [1994(a)]). In fact, the description in Dykema [1993(b)] of D is a special case of the description that follows.

Let us now describe the type I part, D , in (2). (A more precise discussion is found in §7.) Given a faithful state, ψ , on a finite dimensional von Neumann algebra, $D = \bigoplus_{j=1}^K M_{n_j}(\mathbf{C})$, we write

$$(D, \psi) = \bigoplus_{j=1}^K \bigoplus_{\alpha_{j,1}, \dots, \alpha_{j,n_j}} M_{n_j}(\mathbf{C})$$

to mean that the restriction of ψ to the j th summand of D is given by a diagonal density matrix with $\alpha_{j,1}, \dots, \alpha_{j,n_j}$ down the diagonal. If

$$(A_1, \phi_1) = \bigoplus_{j=1}^{K_1} \bigoplus_{\alpha_{j,1}, \dots, \alpha_{j,n_j}} M_{n_j}(\mathbf{C})$$

$$(A_2, \phi_2) = \bigoplus_{j=1}^{K_2} \bigoplus_{\beta_{j,1}, \dots, \beta_{j,m_j}} M_{m_j}(\mathbf{C})$$

and if $\mathcal{M}$ is their free product as in (1), to determine whether $\mathcal{M}$ is a factor and, when it is not, to find the type I part of $\mathcal{M}$, we first “mate” every simple summand of A_1 with every simple summand of A_2 and examine the offspring, if any. When the j th summand of A_1 , namely $\bigoplus_{\alpha_{j,1}, \dots, \alpha_{j,n_j}} M_{n_j}(\mathbf{C})$, is mated with the k th summand of A_2 , namely $\bigoplus_{\beta_{k,1}, \dots, \beta_{k,m_k}} M_{m_k}(\mathbf{C})$, then there can be offspring only if at least one of n_j and m_k is equal to 1, *i.e.* if at least one of the simple summands

is just a copy of $\mathbf{C}$ in the corresponding A_ι . In that case, say if $n_j = 1$, then there is offspring if and only if

$$\sum_{i=1}^{m_k} \frac{1}{\beta_{k,i}} < \frac{1}{1 - \alpha_{j,1}},$$

and then the offspring is

$$M_{m_k}(\mathbf{C}) \quad \text{where} \quad \gamma_i = \beta_{k,i} \left(1 - (1 - \alpha_{j,1}) \sum_{p=1}^{m_j} \frac{1}{\beta_{k,p}} \right).$$

Analogous offspring results if $m_k = 1$ instead of $n_j = 1$. A system of matrix units for the offspring is the meet, $\otimes$, defined in Dykema [1995(a)], of systems of matrix units for the simple summands that were mated, $M_{n_j}(\mathbf{C})$ and $M_{m_k}(\mathbf{C})$, (see §7). Thus the support projection of this offspring lies under the support projection of $M_{n_j}(\mathbf{C})$ and the support projection of $M_{m_k}(\mathbf{C})$. The type I part of $\mathcal{M}$ is equal to the direct sum of all offspring produced in this way by all possible matings of the original simple summands. If none of the matings were successful in producing offspring, then $\mathcal{M}$ is a factor.

For example, we have

$$\left(\begin{smallmatrix} \mathbf{C} & \mathbf{C} \\ \frac{1}{5} & \frac{4}{5} \end{smallmatrix} \right) * \begin{smallmatrix} M_2(\mathbf{C}) \\ \frac{2}{3}, \frac{1}{3} \end{smallmatrix} = \mathcal{M}_a \oplus \begin{smallmatrix} M_2(\mathbf{C}) \\ \frac{1}{15}, \frac{1}{30} \end{smallmatrix} \quad (3a)$$

$$\left(\begin{smallmatrix} \mathbf{C} & \mathbf{C} \\ \frac{1}{4} & \frac{3}{4} \end{smallmatrix} \right) * \begin{smallmatrix} M_2(\mathbf{C}) \\ \frac{2}{3}, \frac{1}{3} \end{smallmatrix} = \mathcal{M}_b \quad (3b)$$

$$\begin{smallmatrix} M_2(\mathbf{C}) \\ \frac{1}{4}, \frac{3}{4} \end{smallmatrix} * \begin{smallmatrix} M_2(\mathbf{C}) \\ \frac{2}{3}, \frac{1}{3} \end{smallmatrix} = \mathcal{M}_c \quad (3c)$$

$$\left(\begin{smallmatrix} M_2(\mathbf{C}) & \mathbf{C} \\ \frac{1}{10(1+\sqrt{8})}, \frac{\sqrt{8}}{10(1+\sqrt{8})} \end{smallmatrix} \oplus \begin{smallmatrix} \mathbf{C} \\ \frac{9}{10} \end{smallmatrix} \right) * \left(\begin{smallmatrix} M_2(\mathbf{C}) & \mathbf{C} \\ \frac{1}{3}, \frac{1}{6} \end{smallmatrix} \oplus \begin{smallmatrix} \mathbf{C} \\ \frac{1}{2} \end{smallmatrix} \right) = \mathcal{M}_d \oplus \begin{smallmatrix} M_2(\mathbf{C}) \\ \frac{1}{30}, \frac{1}{60} \end{smallmatrix} \oplus \begin{smallmatrix} \mathbf{C} \\ \frac{2}{5} \end{smallmatrix} \quad (3d)$$

$$\left(\begin{smallmatrix} M_2(\mathbf{C}) & \mathbf{C} \\ \frac{4}{20}, \frac{1}{20} \end{smallmatrix} \oplus \begin{smallmatrix} \mathbf{C} \\ \frac{3}{4} \end{smallmatrix} \right) * \left(\begin{smallmatrix} M_2(\mathbf{C}) & \mathbf{C} \\ \frac{1}{3}, \frac{1}{6} \end{smallmatrix} \oplus \begin{smallmatrix} \mathbf{C} \\ \frac{1}{2} \end{smallmatrix} \right) = \mathcal{M}_e \oplus \begin{smallmatrix} \mathbf{C} \\ \frac{1}{4} \end{smallmatrix} \quad (3e)$$

$$\left(\begin{smallmatrix} \mathbf{C} & \mathbf{C} \\ \frac{1}{41} & \frac{40}{41} \end{smallmatrix} \right) * \left(\begin{smallmatrix} M_3(\mathbf{C}) & M_2(\mathbf{C}) & M_2(\mathbf{C}) & \mathbf{C} \\ \frac{1}{16}, \frac{1}{16}, \frac{1}{8} & \frac{1}{40}, \frac{9}{40} & \frac{1}{8}, \frac{1}{8} & \frac{1}{4} \end{smallmatrix} \right) = \mathcal{M}_f \oplus \begin{smallmatrix} M_3(\mathbf{C}) \\ \frac{1}{656}, \frac{1}{656}, \frac{1}{328} \end{smallmatrix} \oplus \begin{smallmatrix} M_2(\mathbf{C}) \\ \frac{25}{328}, \frac{25}{328} \end{smallmatrix} \oplus \begin{smallmatrix} \mathbf{C} \\ \frac{37}{164} \end{smallmatrix}, \quad (3f)$$

where $\mathcal{M}_a, \mathcal{M}_b, \mathcal{M}_c, \mathcal{M}_d, \mathcal{M}_e$ and $\mathcal{M}_f$ are type III factors.

Suppose that $\mathcal{M}$ is not a factor, *i.e.* that some matings result in offspring, then one easily sees that there is a *dominant projection*, p , which is by definition a minimal and central projection of A_ι , for $\iota = 1$ or $\iota = 2$, such that all of the offspring are the progeny of the mating of p with simple summands from $A_{\iota'}$ where $\iota' \neq \iota$. For example, in (3a), (3d) and (3f), the only dominant projection is $0 \oplus 1$ in the algebra on the left side of the $*$, whereas in (3f), both $0 \oplus 1$ on the left and $0 \oplus 1$ on the right side of $*$ are dominant projections. The dominant projection $p \in A_\iota$, if it exists, must be the largest (with respect to ϕ_ι) of the projections in A_ι that are both minimal and central.

We now turn to a discussion of the type III factor, $\mathcal{M}_0$, appearing in Theorem 1. Connes [1973] defined a state, ϕ , on a von Neumann algebra, $\mathcal{M}$, to be almost periodic if $L^2(\mathcal{M}, \phi)$ has a basis of eigenvectors for the

modular operator, Δ_ϕ , of ϕ . In that case, Arveson–Connes spectral theory (Arveson [1974], Connes [1973]) provides a decomposition of $\mathcal{M}$ as a direct sum of spectral subspaces. If, in addition, the centralizer, $\mathcal{M}_\phi$, of ϕ is a factor, then, it is easily seen, the point spectrum of Δ_ϕ is a subgroup of $\mathbf{R}_+^*$ and every spectral subspace is of the form $\mathcal{M}_\phi v$ or $v^* \mathcal{M}_\phi$ for an isometry v . In analogy with Connes’ results about full factors, Connes [1974], we call an almost periodic state, ϕ , having centralizer that is a factor an *extremal almost periodic* state, and we define $Sd(\phi)$ to be the point spectrum of Δ_ϕ . (At the end of §5, we show that term “extremal” is in some sense appropriate.) By Corollary 3.2.7 of Connes [1973], from the fact that $\mathcal{M}_\phi$ is a factor we have also that Connes’ invariant $S(\mathcal{M})$ is equal to the closure of $Sd(\phi)$. Hence in Theorem 1, ϕ_0 is extremal almost periodic and $Sd(\phi_0)$ is found from the initial data, namely ϕ_1 and ϕ_2 . Moreover, regarding Connes’ classification of $\mathcal{M}_0$ as type III_λ , we have: $\mathcal{M}_0$ is type III_λ for $0 < \lambda < 1$ if $Sd(\phi_0) = \lambda^{\mathbf{Z}} = \{\lambda^n \mid n \in \mathbf{Z}\}$ and otherwise $Sd(\phi_0)$ is dense in $\mathbf{R}_+^*$, and $\mathcal{M}_0$ must be type III_1 .

For example,

in (3a), $\mathcal{M}_a$ is a type III_λ factor for $\lambda = 1/2$;

in (3b), $\mathcal{M}_b$ is a type III_λ factor for $\lambda = 1/2$;

in (3c), $\mathcal{M}_c$ is a type III_1 factor and the free product state, ϕ , has $Sd(\phi)$ equal to the (dense) subgroup of $\mathbf{R}_+^*$ generated by $1/3$ and $1/2$;

in (3d), $\mathcal{M}_d$ is a type III_λ factor for $\lambda = 1/\sqrt{2}$;

in (3e), $\mathcal{M}_e$ is a type III_λ factor for $\lambda = 1/2$;

in (3f), $\mathcal{M}_f$ is a type III_1 factor and the free product state, ϕ , has $Sd(\phi)$ equal to the (dense) subgroup of $\mathbf{R}_+^*$ generated by $1/2$ and $1/9$.

Extremal almost periodic states play a prominent role in the proof of Theorem 1. An example of the sort of results proved and used in the course of this paper is the following theorem.

Theorem 2 *Suppose for $\iota = 1, 2$ that A_ι is a separable type III factor and ϕ_ι is an extremal almost periodic state on A_ι whose centralizer is isomorphic to the hyperfinite II_1 factor, R , or to $L(F_\infty)$. Let*

$$(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2).$$

Then $\mathcal{M}$ is a type III factor, ϕ is extremal almost periodic and $Sd(\phi)$ is the subgroup of $\mathbf{R}_+^*$ generated by $Sd(\phi_1) \cup Sd(\phi_2)$.

By taking inductive limits, Theorem 1 can be extended to more general algebras.

Theorem 3 *Let*

$$(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2),$$

where ϕ_1 and ϕ_2 are faithful states, at least one of which is not a trace, where A_1 and A_2 are nontrivial algebras, each of which is one of the following:

- (i) *finite dimensional*
- (ii) *$B(\mathcal{H})$ for separable, infinite dimensional Hilbert space $\mathcal{H}$*
- (iii) *a type III factor on which ϕ_ι is extremal almost periodic and has centralizer isomorphic to the hyperfinite II_1 factor, R , or to $L(F_\infty)$;*

- (iv) a diffuse hyperfinite von Neumann algebra on which ϕ_ι is a trace
- (v) an interpolated free group factor (of type II_1), see Rădulescu [1994], Dykema [1994(a)]
- (vi) a possibly infinite direct sum of algebras from (i)-(v).

Then

$$\mathcal{M} = \mathcal{M}_0 \oplus D \quad \text{or} \quad \mathcal{M} = \mathcal{M}_0,$$

with $\mathcal{M}_0$ and D as described in Theorem 1, (and more explicitly below).

In Theorem 3, one finds the finite dimensional part, D , of $\mathcal{M}$, if it exists, by mating every matrix algebra direct summand of A_1 with every matrix algebra summand of A_2 , as described above, and D is the sum of the offspring of these matings.

Theorem 4 *Let*

$$(\mathcal{M}, \phi) = \bigstar_{\iota \in I} (A_\iota, \phi_\iota),$$

where I is finite or countably infinite, $|I| \geq 2$, each ϕ_ι is a faithful state on A_ι , at least one ϕ_ι is not a trace and each A_ι is of the form described in Theorem 3. Then

$$\mathcal{M} = \mathcal{M}_0 \oplus D \quad \text{or} \quad \mathcal{M} = \mathcal{M}_0,$$

with $\mathcal{M}_0$ and D as described in Theorem 1, (and more explicitly below).

In this case, one finds the finite dimensional part, D , of $\mathcal{M}$, if it exists, by, for every choice of a matrix algebra direct summand, say $M_{n_\iota}(\mathbf{C})$ from each A_ι , mating them all together, and letting D be the sum of the offspring resulting from all such matings. A mating of one summand from each A_ι results in offspring only if all but one of the n_ι equal 1, say $n_\iota = 1$ if $\iota \in I \setminus \{\iota'\}$, and then offspring results if and only if

$$\frac{1}{\sum_{\iota \in I \setminus \{\iota'\}} (1 - \alpha_{\iota,1})} > \sum_{j=1}^{n_{\iota'}} \frac{1}{\alpha_{\iota',j}}$$

and the offspring is

$$M_{n_{\iota'}}(\mathbf{C}) \quad \text{where} \quad \gamma_i = \alpha_{\iota',i} \left(1 - \left(\sum_{\iota \in I \setminus \{\iota'\}} (1 - \alpha_{\iota,1}) \right) \sum_{p=1}^{n_{\iota'}} \frac{1}{\alpha_{\iota',p}} \right).$$

We also have, as in Theorem 1, that $\mathcal{M}_0$ is a type III factor, $\phi_0 = \phi|_{\mathcal{M}_0}$ is extremal almost periodic and $Sd(\phi_0)$ is equal to the subgroup of $\mathbf{R}_+^*$ generated by the union over $\iota \in I$ of the point spectra of Δ_{ϕ_ι} .

For example,

$$(B(\mathcal{H}), \text{Tr} \left(\begin{pmatrix} 1/2 & * & & \\ & 1/4 & & \\ & & 1/8 & \\ & & & \ddots \end{pmatrix} \cdot \right)) * (B(\mathcal{H}), \text{Tr} \left(\begin{pmatrix} 3/4 & & & \\ & 3/16 & & \\ & & 3/64 & \\ & & & \ddots \end{pmatrix} \cdot \right)) = \mathcal{M}_g \quad (4g)$$

$$\left((B(\mathcal{H}), \text{Tr} \left(\begin{pmatrix} 1/2 & & & \\ & 1/4 & & \\ & & 1/8 & \\ & & & \ddots \end{pmatrix} \cdot \right)) \oplus_{9/10} \mathbf{C} \right) *_{\frac{3}{4}, \frac{1}{4}} M_2(\mathbf{C}) = \mathcal{M}_h \oplus_{\frac{7}{20}, \frac{7}{60}} M_2(\mathbf{C}) \quad (4h)$$

$$\bigstar_{n \geq 1} M_2(\mathbf{C}) = \mathcal{M}_k, \quad (4k)$$

where $\mathcal{M}_g$, $\mathcal{M}_h$ and $\mathcal{M}_k$ are type III factors and

in (4g), $\mathcal{M}_g$ is a type III_λ factor for $\lambda = 1/2$;

in (4h), $\mathcal{M}_h$ is a type III_1 factor and the free product state restricted to $\mathcal{M}_h$, call it ϕ_0 , has $Sd(\phi_0)$ equal to the subgroup of $\mathbf{R}_+^*$ generated by $1/2$ and $1/3$;

in (4k), $\mathcal{M}_k$ is a type III_1 factor and the free product state, ϕ , has $Sd(\phi)$ equal to the subgroup of $\mathbf{R}_+^*$ generated by $\{\sqrt[n]{2} \mid n \geq 1\}$.

The proofs of the above theorems involve what is loosely speaking a “type III version” of the following fact from group theory. Let $F_2 = \langle a, b \rangle$ be the free group with free generators a and b . Let $\pi : F_2 \rightarrow \mathbf{Z}$ be the group homomorphism such that $\pi(a) = 0$ and $\pi(b) = 1$. Then $\ker \pi \cong F_\infty$ and $\ker \pi$ is freely generated by $(b^n a b^{-n})_{n \in \mathbf{Z}}$. By “type III version” we mean that trying to find the centralizer of certain free product states is like trying to find $\ker \pi$ above, where instead of b we have a nonunitary isometry. Of course, there are some technically more involved aspects, but the basic idea is as easy as this. We plan to write up the proof of a special case, namely $M_2(\mathbf{C}) * M_2(\mathbf{C})$ with respect to arbitrary faithful states, in Dykema [SNU]. This proof will contain most of the essential ideas of the proof of the general case, but should more transparent, and we would recommend it to those interested in an introduction to the techniques.

Many results of this paper are proved using the technique of *free etymology*. This technique consists of proving the freeness of certain algebras by looking at words in those algebras, and investigating the “roots” of those words. To be more specific, suppose we want to show that a family $(A_\iota)_{\iota \in I}$ of subalgebras of A is free in (A, ϕ) , and each A_ι is defined in terms of some other subalgebras $(B_j)_{j \in J}$ that we know are free. Since we work so much with words, let us use a special notation, which we have used before in Dykema [1993(b)], Dykema [1994(a)] and Dykema [1995(a)].

Definition 5 Suppose A is an algebra, $\phi : A \rightarrow \mathbf{C}$ is linear. For $B \subseteq A$ any subalgebra, we define B° (pronounced B -bubble) to be $B^\circ = B \cap \ker \phi$. Suppose $X_\iota \subseteq A$, are subsets, $(\iota \in I)$. The set of *reduced words* in $(X_\iota)_{\iota \in I}$ is denoted $\Lambda^\circ((X_\iota)_{\iota \in I})$, and is defined to be the set of $a_1 a_2 \cdots a_n$ such that $n \geq 1$, $a_j \in X_{\iota_j}$ and $\iota_1 \neq \iota_2 \neq \cdots \iota_n$. If $|I| = 2$, we may call them *alternating products*.

Now a free etymology proof of the freeness of $(A_\iota)_{\iota \in I}$ would go as follows: we need only show $\phi(x) = 0$ for every $x \in \Lambda^\circ((A_\iota)_{\iota \in I})$; after doing some work, one shows that x is equal to, for example if A is a von Neumann algebra, a strong-operator limit of sums of reduced words in $(B_j^\circ)_{j \in J}$, and hence one concludes that $\phi(x) = 0$.

In §8, using a result of Barnett [1995], we show in many cases that $\mathcal{M}_0$ in Theorems 1, 3 and 4 is a full factor. Thus, for these factors, Connes’ invariant for full factors, $Sd(\mathcal{M}_0)$, is equal to $Sd(\phi_0)$, and thereby, for instance by taking free products as in example (4k), we obtain for an arbitrary countable dense subgroup, Γ , of $\mathbf{R}_+^*$, a free product factor that is a full type III_1 factor, has Γ as its Sd invariant and has extremal almost periodic state with centralizer isomorphic to $L(F_\infty)$. See Corollary 4.4 of Connes [1974] and Golodets and Nessonov [1987] for other constructions of full III_1 factors with arbitrary Sd invariant.

Acknowledgements. Most of this work was done while I was at the Fields Institute during the special year in Operator Algebras, 1994/95. I would like to thank George Elliott for the organizing that marvelous year.

Contents:

§1. Free Subcomplementation	p.7
§2. Freeness with amalgamation over a subalgebra	p.11
§3. Conditional Expectations	p.11
§4. Extremal almost periodic states	p.13
§5. Some technical lemmas	p.16
§6. Free products of factors with respect to extremal almost periodic states.	p.19
§7. Free products of finite dimensional algebras	p.31
§8. Fullness of free product factors	p.44
§9. Questions	p.45
References	p.46

1. Free Subcomplementation.

In Dykema [1993(b)], the free product of arbitrary finite dimensional or AFD von Neumann algebras with respect to faithful traces was determined, and found to be an interpolated free group factor (Rădulescu [1994], Dykema [1994(a)]) possibly direct sum a finite dimensional algebra. The free product of finite dimensional algebras was determined using the technique that is now called *free etymology*, and passing to AFD algebras was permitted by the use of the notion of a *standard embedding* of interpolated free group factors. The definition of a standard embedding of interpolated free group factors, $L(F_s) \hookrightarrow L(F_t)$ is a generalization of the sort of embedding $L(F_n) \hookrightarrow L(F_m)$, $n < m$ or $n = m = \infty$, obtained by mapping the n free generators of F_n to n (not all) of the m free generators of F_m .

The characterization of standard embeddings in Proposition 1.2 is pleasing to the intuition and will be used in this paper.

Definition 1.1 Suppose $(\mathcal{M}, \phi)$ is a von Neumann algebra with normal state, and let $1 \in A \subseteq \mathcal{M}$ be a von Neumann subalgebra. We say that A is *freely complemented* in $(\mathcal{M}, \phi)$ if there is a von Neumann subalgebra $B \subseteq \mathcal{M}$ such that $\mathcal{M}$ is generated by $A \cup B$ and A and B are free in $(\mathcal{M}, \phi)$. To be more specific, we can say that A is freely complemented by B .

Proposition 1.2 Suppose $\pi : L(F_s) \hookrightarrow L(F_t)$ is an injective, normal, trace-preserving $*$ -homomorphism. Then π is a standard embedding if and only if $\pi(L(F_s))$ is freely complemented in $L(F_t)$ by an algebra isomorphic to $L(F_r) \oplus \mathbf{C}$ or simply to $L(F_r)$, some $1 < r \leq \infty$.

Proof Let $A = \pi(L(F_s))$ be freely complemented in $L(F_t)$ by $B \cong L(F_r) \oplus \mathbf{C}$ or $\cong L(F_r)$. Then by 4.4 and 1.2 of Dykema [1993(b)], π is a standard embedding. Since for any two standard embeddings $\pi_1 : L(F_s) \hookrightarrow L(F_t)$ and $\pi_2 : L(F_s) \hookrightarrow L(F_t)$, there is an automorphism α of $L(F_t)$ such that $\pi_1 = \pi_2 \circ \alpha$, (this is easily seen from the definition of a standard embedding Dykema [1993(b)]), and since for any s and t we can find $B = L(F_r) \oplus \mathbf{C}$ and a trace on it such that $L(F_s) * B \cong L(F_t)$, it follows that every standard embedding is of this form. $\square$

The above characterization of standard embeddings allows one to easily prove the following useful facts about standard embeddings, some of which first appeared in Dykema [1993(b)].

Proposition 1.3

- (i) *The composition of standard embeddings is a standard embedding.*
- (ii) *Suppose $A_n = L(F_{t_n})$, $1 < t_n \leq \infty$ and $\pi_n : A_n \rightarrow A_{n+1}$ is a standard embedding ($n \geq 1$). Consider the inductive limit von Neumann algebra (taken with respect to the unique traces on A_n), $A = \varinjlim (A_n, \pi_n)$ with the associated inclusions $\psi_n : A_n \hookrightarrow A$. Then $A = L(F_t)$ where $t = \lim_{n \rightarrow \infty} t_n$ and each ψ_n is a standard embedding.*

We now turn to an extension of the notion of a freely complemented subalgebra.

Definition 1.4 Suppose $(\mathcal{M}, \phi)$ is a von Neumann algebra with normal state, and let $1 \in A \subseteq \mathcal{M}$ be a von Neumann subalgebra. Let A_ϕ denote the centralizer of $\phi|_A$. We say that A is *freely subcomplemented* in $(\mathcal{M}, \phi)$ if, for some index set I , there is for each $\iota \in I$ a self-adjoint projection $h_\iota \in A_\phi$ and a von Neumann subalgebra $B_\iota \subseteq h_\iota \mathcal{M} h_\iota$ such that

- (i) $\mathcal{M} = W^*(A \cup \bigcup_{\iota \in I} B_\iota)$
- (ii) B_ι and $h_\iota \mathcal{M}_{I \setminus \{\iota\}} h_\iota$ are free in $(h_\iota \mathcal{M} h_\iota, \phi(h_\iota)^{-1} \phi|_{h_\iota \mathcal{M} h_\iota})$, where $\mathcal{M}_{I \setminus \{\iota_0\}} \stackrel{\text{def}}{=} W^*(A \cup \bigcup_{\iota \in I \setminus \{\iota_0\}} B_\iota)$.

To be more precise, we will say that A is freely subcomplemented in $(\mathcal{M}, \phi)$ by $(B_\iota)_{\iota \in I}$ and with associated projections $(h_\iota)_{\iota \in I}$.

The condition (ii)' of the following lemma will be the most useful one for proving properties of free subcomplementation.

Lemma 1.5 *Suppose $(\mathcal{M}, \phi)$, A , h_ι and B_ι are as in Definition 1.4, except without assuming (ii) holds and suppose also that the GNS representation of $\mathcal{M}$ associated to ϕ is faithful. Let Ω be the set of reduced words $a_1 a_2 \cdots a_n \in \Lambda^\circ(A, \bigcup_{\iota \in I} B_\iota^\circ)$ such that whenever $2 \leq j \leq n-1$, $a_j \in A$, and $a_{j-1}, a_{j+1} \in B_\iota^\circ$ for some $\iota \in I$, then $a_j \in (h_\iota A h_\iota)^\circ$. Then $\text{span}(\Omega \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{M}$. Consider the condition*

- (ii)' $\phi(z) = 0 \ \forall z \in \Omega \setminus A$.

Then (ii)' $\implies$ (ii). Moreover, (ii)' implies the existence of a s.o.-continuous projection of norm 1, $E : \mathcal{M} \rightarrow A$, such that $E(z) = 0 \ \forall z \in \Omega \setminus A$. Conversely, if each h_ι has central support in A_ϕ equal to 1 then (ii) $\implies$ (ii)'.

Proof One easily sees that $\text{span}(\Omega \cup \{1\})$ is the $*$ -algebra generated by $A \cup \bigcup_{\iota \in I} B_\iota$, hence is s.o.-dense in $\mathcal{M}$. For $F \subseteq I$, let Ω_F be the set of all words $z \in \Omega$ such that whenever z has a letter coming from B_ι° , $\iota \in I$, then $\iota \in F$. Let us show that (ii)' implies the existence of the projection $E : \mathcal{M} \rightarrow A$. We will realize this projection by compressing to $L^2(A, \phi)$ in the Hilbert space $L^2(\mathcal{M}, \phi)$. The defining mapping $\mathcal{M} \rightarrow L^2(\mathcal{M}, \phi)$ will be denoted $x \mapsto \hat{x}$ and we let $\xi \stackrel{\text{def}}{=} \hat{1}$.

Lemma 1.6 *If $\mathcal{P}$ is a von Neumann algebra with normal state ϕ and if $h, k \in \mathcal{P}_\phi$ are self-adjoint projections in the centralizer of ϕ such that $\phi(hk) = 0$, then $\phi(hak) = 0 \ \forall a \in \mathcal{P}$.*

Proof $0 = \phi(hk) = \phi(khk)$. The left support of kh is the support of khk which is $k - k \wedge (1 - h)$. Since $khk \geq 0$ and $\phi(khk) = 0$ we have $\phi(k - k \wedge (1 - h)) = 0$. Thus

$$\phi(hak) = \phi(khak) = \phi((k - k \wedge (1 - h))ha(k - k \wedge (1 - h))) = 0.$$

Hence Lemma 1.6 is proved. $\square$

Continuing with the proof of Lemma 1.5, using Lemma 1.6 we see for $\iota_1, \iota_2 \in I$ that

$$\begin{aligned} h_{\iota_1}Ah_{\iota_2} &\subseteq \ker \phi && \text{if } \phi(h_{\iota_1}h_{\iota_2}) = 0 \\ h_{\iota_1}Ah_{\iota_2} &= (h_{\iota_1}Ah_{\iota_2}) \cap \ker \phi + \mathbf{C}h_{\iota_1}h_{\iota_2} && \text{if } \phi(h_{\iota_1}h_{\iota_2}) \neq 0 \end{aligned} \quad (5)$$

Using (ii)', equation (5) and that $\text{span}(\Omega \cup \{1\})$ is dense in $\mathcal{M}$, we find that $L^2(\mathcal{M}, \phi)$ can be characterized as follows.

Let $\alpha \notin I$, $\mathcal{H}_\alpha \stackrel{\text{def}}{=} L^2(A, \phi|_A) \ominus \mathbf{C}\hat{1}$, for $\iota \in I$ let $\mathcal{H}_\iota \stackrel{\text{def}}{=} L^2(B_\iota, \phi|_{B_\iota}) \ominus \mathbf{C}\hat{h}_\iota$, and let

$$\mathcal{H} \stackrel{\text{def}}{=} \mathbf{C}\xi \oplus \bigoplus_{\substack{n \geq 1 \\ j_k \in I \cup \{\alpha\} \\ j_1 \neq j_2 \neq \dots \neq j_n}} \mathcal{H}_{j_1} \otimes \mathcal{H}_{j_2} \otimes \dots \otimes \mathcal{H}_{j_n}.$$

Then $L^2(\mathcal{M}, \phi)$ is the closed subspace of $\mathcal{H}$ that is spanned by ξ together with all vectors

$$\zeta_1 \otimes \dots \otimes \zeta_n \in \mathcal{H}_{j_1} \otimes \dots \otimes \mathcal{H}_{j_n}, \quad j_1 \neq \dots \neq j_n$$

such that

$$\begin{aligned} \text{if } 2 \leq k \leq n-1, j_k = \alpha, \text{ (hence } j_{k-1}, j_{k+1} \in I), \text{ then } \zeta_k &\in \overline{\{\hat{a} \mid a \in h_{j_{k-1}}Ah_{j_{k+1}} \cap \ker \phi\}} \\ \text{if } n \geq 2 \text{ and } j_1 = \alpha, \text{ (hence } j_2 \in I), \text{ then } \zeta_1 &\in \overline{\{\hat{a} \mid a \in Ah_{j_2} \cap \ker \phi\}} \\ \text{if } n \geq 2 \text{ and } j_n = \alpha, \text{ (hence } j_{n-1} \in I), \text{ then } \zeta_n &\in \overline{\{\hat{a} \mid a \in h_{j_{n-1}}A \cap \ker \phi\}}. \end{aligned}$$

Let P_A be the self-adjoint projection of $L^2(\mathcal{M}, \phi)$ onto $L^2(A, \phi|_A) = \mathbf{C}\xi \oplus \mathcal{H}_\alpha \subseteq L^2(\mathcal{M}, \phi)$. Let $E : B(L^2(\mathcal{M}, \phi)) \rightarrow B(L^2(A, \phi|_A))$ be defined by $E(x) = P_A x P_A$. Then E is a projection of norm 1 and is s.o.-continuous. We easily see that

$$\begin{aligned} E(a) &= a \quad \forall a \in A \\ E(z) &= 0 \quad \forall z \in \Omega \setminus A. \end{aligned}$$

Thus, by s.o.-continuity of E , we see that E projects $\mathcal{M}$ onto A .

Now using the projection E , we will show (ii)' $\implies$ (ii). Let $\iota \in I$. We will show that B_ι and $h_\iota \mathcal{M}_{I \setminus \{\iota\}} h_\iota$ are free in $h_\iota \mathcal{M} h_\iota$. Clearly $\text{span}(\Omega_{I \setminus \{\iota\}} \cup \{1\})$ is a dense $*$ -subalgebra of $\mathcal{M}_{I \setminus \{\iota\}}$. Using the projection $E : \mathcal{M} \rightarrow A$, we see that each element of $(h_\iota \mathcal{M}_{I \setminus \{\iota\}} h_\iota)^\circ$ is the s.o.-limit of a bounded net in $\text{span}(h_\iota(\Omega_{I \setminus \{\iota\}} \setminus A)h_\iota \cup (h_\iota Ah_\iota)^\circ)$, hence it will suffice to show $\phi(y) = 0$ whenever

$$y \in \Lambda^\circ(h_\iota(\Omega_{I \setminus \{\iota\}} \setminus A)h_\iota \cup (h_\iota Ah_\iota)^\circ, B_\iota^\circ).$$

But by regrouping we see that $y \in (\Omega \setminus A) \cup (h_\iota Ah_\iota)^\circ$, hence $\phi(y) = 0$ by (ii)'.

Assume that each h_ι has central support in A_ϕ equal to 1 and let us show (ii) $\implies$ (ii)'. Take any $z = a_1 a_2 \dots a_n \in \Omega_F \setminus A$, where F is a finite subset of I . It will suffice to show $\phi(z) = 0$, and we will proceed by induction on $|F|$,

proving the $|F| = 1$ case and the inductive step simultaneously. Let $\iota \in F$ be such that z has a letter from B_ι° . Now we may assume $n \geq 3$ and $a_1, a_n \in A$. By hypothesis, there are partial isometries $v_0, \dots, v_k \in A_\phi$ such that $v_j^* v_j \leq h_\iota \forall 1 \leq j \leq k$ and $\sum_{j=1}^k v_j v_j^* = 1$. Now $\phi(z) = \sum_{j_1, j_2=0}^k \phi(v_{j_1} v_{j_1}^* z v_{j_2} v_{j_2}^*) = \sum_{j=0}^k \phi(v_j^* z v_j)$, so we may assume without loss of generality that $a_1 \in h_\iota A$ and $a_n \in A h_\iota$. Regrouping in z by singling out all letters from B_ι° and grouping all other strings of neighboring letters together, we see that $z \in \Lambda^\circ(h_\iota \Omega_{F \setminus \{\iota\}} h_\iota, B_\iota^\circ)$ and if a letter of z is in $h_\iota \Omega_{F \setminus \{\iota\}} h_\iota \cap A$ then it is in $(h_\iota A h_\iota)^\circ$, unless it is the first or the last letter, in which case it is in $h_\iota A h_\iota$. Using if necessary $h_\iota A h_\iota = (h_\iota A h_\iota)^\circ + \mathbf{C} h_\iota$ to expand the first and/or the last letter, we see that z is a linear combination of up to four elements of

$$\Lambda^\circ((h_\iota \Omega_{F \setminus \{\iota\}} h_\iota \setminus A) \cup (h_\iota A h_\iota)^\circ, B_\iota^\circ).$$

By inductive hypothesis in the case $|F| > 1$ and by the fact that $\Omega_{F \setminus \{\iota\}} \subseteq A^\circ$ if $|F| = 1$, we have that

$$z \in \text{span} \Lambda^\circ((h_\iota \mathcal{M}_{I \setminus \{\iota\}} h_\iota)^\circ, B_\iota^\circ),$$

hence by (ii), $\phi(z) = 0$. □

Now we show that the composition of (certain) freely subcomplemented embeddings is a freely subcomplemented embedding.

Corollary 1.7 *Suppose*

$$1 \in A_1 \subseteq A_2 \subseteq \dots \subseteq \mathcal{M} \tag{6}$$

is a finite or infinite chain of inclusions of von Neumann algebras, ϕ is a normal state on $\mathcal{M}$ and

$$\mathcal{M} = \begin{cases} A_n & \text{if the chain (6) is finite of length } n \\ \bigcup_{n \geq 1} A_n & \text{if the chain (6) is infinite} \end{cases}$$

Suppose for all j that A_j is freely subcomplemented in $(A_{j+1}, \phi|_{A_{j+1}})$ by $(B_\iota)_{\iota \in I_j}$ with associated projections $(h_\iota)_{\iota \in I_j}$ such that each h_ι has central support in $(A_j)_\phi$ equal to 1 and is Murray–von Neumann equivalent in $(A_j)_\phi$ to a projection in A_1 . Then A_1 is freely subcomplemented in $\mathcal{M}$ by (algebras isomorphic to) $(B_\iota)_{\iota \in I}$, with $I = \bigcup_j I_j$.

The above result is a special case of the next one.

Corollary 1.8 *Let J be a set and $\leq$ a well-ordering on J . Suppose $\forall j \in J$ A_j is a von Neumann algebra with normal state ϕ_j and suppose there are compatible, ϕ_j –preserving inclusions $A_{j_1} \subseteq A_{j_2} \forall j_1 \leq j_2$. Let $(\mathcal{M}, \phi) = \varinjlim_{j \in J} (A_j, \phi_j)$ be the inductive limit von Neumann algebra. Let $j_0 \in J$ denote the smallest element and suppose $\forall j \in J \setminus \{j_0\}$ that $\mathcal{M}_{\leq j} \stackrel{\text{def}}{=} W^*(\{1\} \cup \bigcup_{j_1 \leq j} A_{j_1})$ is freely subcomplemented in $W^*(\mathcal{M}_{\leq j} \cup A_j)$ (with respect to the restriction of ϕ to $W^*(\mathcal{M}_{\leq j} \cup A_j)$) by $(B_\iota)_{\iota \in I_j}$ with associated projections $(h_\iota)_{\iota \in I_j}$, such that each h_ι has central support in $(A_j)_\phi$ equal to 1 and is Murray–von Neumann equivalent to a projection in A_{j_0} . Then A_{j_0} is freely subcomplemented in $(\mathcal{M}, \phi)$. by (algebras isomorphic to) $(B_\iota)_{\iota \in I}$ where $I = \bigcup_{j \in J} I_j$.*

Proof Let $(h_\iota)_{\iota \in I_j}$ be the projections associated to $(B_\iota)_{\iota \in I_j}$. We may suppose $h_\iota \in (A_{j_0})_\phi \forall \iota \in I$. Now it will suffice to show (ii)’ of Lemma 1.5, (with $A = A_{j_0}$ here). Let $z \in \Omega_F \setminus A$, where $F \subseteq I$ is any finite subset and let $j \in J$ be the largest element of J such that $F \cap I_j$ is nonempty. We show $\phi(z) = 0$ by transfinite induction on j . If $j = j_0$ then $\Omega_F \subseteq A$ and there is nothing to prove. Otherwise, regrouping in z by grouping letters from all B_ι° for $\iota \notin I_j$

and A together, we see that z equals an element of $\Lambda^\circ(\Omega_{F \setminus I_j}, \bigcup_{\iota \in I_j} B_\iota^\circ)$ whose every letter lying between letters from B_ι° , $\iota \in I_j$ belongs to $(h_\iota \Omega_{F \setminus I_j} h_\iota \setminus A) \cup (h_\iota A h_\iota)^\circ$. This last set is, by inductive hypothesis, a subset of $(h_\iota \mathcal{M}_{\not\subseteq j} h_\iota)^\circ$. Applying Lemma 1.5(ii)' to the situation of $\mathcal{M}_{\not\subseteq j}$ freely subcomplemented in A_j , we have that $\phi(z) = 0$. $\square$

2. Freeness with amalgamation over a subalgebra.

Freeness of C^* -algebras over a subalgebras was defined by Voiculescu, and this notion of freeness corresponds to his construction of free products with amalgamation over common subalgebras.

Definition 2.1(Voiculescu [1985]) Suppose A is an algebra, $B \subseteq A$ a subalgebra, $E : A \rightarrow B$ a B - B bimodule map. Given subalgebras $B \subseteq A_\iota \subseteq A$ ($\iota \in I$), we say that the family $(A_\iota)_{\iota \in I}$ is *free over B* if $E(a_1 a_2 \cdots a_n) = 0$ whenever $a_j \in A_{\iota_j}$, $\iota_k \neq \iota_{k+1}$ ($1 \leq k \leq n-1$) and $E(a_j) = 0 \ \forall j$. To be emphatic, we will sometimes say the family $(A_\iota)_{\iota \in I}$ is *free with amalgamation over B* .

Proposition 2.2 Suppose A is a C^* -algebra with state ϕ , B is a C^* -subalgebra of A such that the G.N.S. representation associated to $\phi|_B$ is faithful on B , $E : A \rightarrow B$ is a projection of norm one (hence a conditional expectation) from A onto B such that $\phi \circ E = \phi$. Suppose $B \subseteq A_\iota \subseteq A$ are subalgebras. Then to show that $(A_\iota)_{\iota \in I}$ is free with amalgamation over B , it suffices to show that $\phi(a_1 a_2 \cdots a_n) = 0$ whenever $a_j \in A_{\iota_j}$, $\iota_1 \neq \iota_2 \neq \cdots \neq \iota_n$ and $E(a_j) = 0 \ \forall j$.

Proof From the fact that $E(a_j) = 0$ we have, for any $b_1, b_2 \in B$, that $E(a_n b_2) = E(a_n) b_2 = 0 = E(b_1 a_1)$ and hence, if the condition of the proposition holds, we have $0 = \phi((b_1 a_1) a_2 \cdots (a_n b_2)) = \phi \circ E((b_1 a_1) a_2 \cdots (a_n b_2)) = \phi(b_1 E(a_1 \cdots a_n) b_2)$ Letting $(\pi, \mathcal{H}, \xi) = \text{GNS}(B, \phi_B)$, it follows that $0 = \phi(b_1 E(a_1 \cdots a_n) b_2) = \langle \pi(E(a_1 \cdots a_n)) \hat{b}_2, (b_1^*)^\wedge \rangle$ so $\pi(E(a_1 \cdots a_n)) = 0$ and hence $E(a_1 \cdots a_n) = 0$. Thus, the condition of the proposition implies freeness over B . $\square$

Proposition 2.3 Suppose $\mathcal{M}$ is a von Neumann algebra with normal state ϕ , $1 \in A \subseteq \mathcal{M}$ a von Neumann subalgebra, $E = E_A^\mathcal{M} : \mathcal{M} \rightarrow A$ a s.o.-continuous projection of norm 1 such that $\phi \circ E = \phi$. Suppose $A \subseteq \mathcal{N}_j \subseteq \mathcal{M}$ is a von Neumann subalgebra ($j \in J$) such that $\bigcup_{j \in J} \mathcal{N}_j$ generates $\mathcal{M}$ and $(\mathcal{N}_j)_{j \in J}$ is free over A in $(\mathcal{M}, E)$. Suppose for each $j \in J$ A is freely subcomplemented in $\mathcal{N}_j$ by $(B_\iota)_{\iota \in I_j}$ with associated projections $(h_\iota)_{\iota \in I_j}$ such that the central support of each h_ι in the centralizer of $\phi|_A$ is 1. Let $I = \bigcup_{j \in J} I_j$ be a disjoint union. Then A is freely subcomplemented in $\mathcal{M}$ by $(B_\iota)_{\iota \in I}$ with associated projections $(h_\iota)_{\iota \in I}$ and $\mathcal{N}_j$ is freely subcomplemented in $\mathcal{M}$ by $(B_\iota)_{\iota \in I \setminus I_j}$ with associated projections $(h_\iota)_{\iota \in I \setminus I_j}$.

Proof To prove that both A and $\mathcal{N}_j$ are freely subcomplemented in $\mathcal{M}$, using Definition 1.4(ii) and Lemma 1.5(ii)' we see that it will suffice to show $\phi(z) = 0 \ \forall z \in \Omega \setminus A$, with Ω as in Lemma 1.5. However, by regrouping, we see that $z \in \Lambda^\circ((\Omega_{I_j} \setminus A)_{j \in J})$. Using that $\phi(y) = 0 \ \forall y \in \Omega_{I_j} \setminus A$, we see that $\Omega_{I_j} \setminus A \subseteq \mathcal{N}_j \cap \ker E_A^\mathcal{M}$, hence $\phi(z) = 0$. $\square$

3. Conditional Expectations.

Recall Tomiyama's famous result, Tomiyama [1957], that a projection of norm 1, E , from a C^* -algebra A onto a C^* -subalgebra B is necessarily positive and a conditional expectation, i.e. $E(b_1 a b_2) = b_1 E(a) b_2$ for all $a \in A$,

$b_1, b_2 \in B$. The following two lemmas, (which we do not claim are new), form a characterization of conditional expectations in terms of projections on the Hilbert space of a G.N.S. representation.

Lemma 3.1 *Let A be a von Neumann algebra, $B \subseteq A$ a von Neumann subalgebra and ϕ a normal, faithful state on A . Let $(\pi, \mathcal{H}, \xi) = \text{GNS}(A, \phi)$, so $\mathcal{H}$ contains $\{\hat{a} \mid a \in A\}$ as a dense subspace with inner product $\langle \hat{a}_1, \hat{a}_2 \rangle = \phi(a_2^* a_1)$. Let $\mathcal{H}_B = \overline{\{\hat{b} \mid b \in B\}} \subseteq \mathcal{H}$, and let $P : \mathcal{H} \rightarrow \mathcal{H}_B$ be the orthogonal projection. Suppose that $\mathfrak{A}$ is a σ -weakly dense subset of A , and that $P\pi(a)|_{\mathcal{H}_B} \in P\pi(B)|_{\mathcal{H}_B}$ for every $a \in \mathfrak{A}$. Then there is a projection of norm 1, $E : A \rightarrow B$, such that $\phi \circ E = \phi$. Moreover, E is continuous from the s.o.-topology on A (acting on $\mathcal{H}$) to the s.o.-topology on B (acting on $\mathcal{H}_B$).*

Proof We have that π is a normal, faithful representation of A on $\mathcal{H}$ and $\pi_B : B \rightarrow B(\mathcal{H}_B)$ given by $\pi_B(b) = P\pi(b)|_{\mathcal{H}_B}$ is a normal, faithful representation of B on $\mathcal{H}_B$, and in fact $(\pi_B, \mathcal{H}_B, \xi) = \text{GNS}(B, \phi|_B)$. Hence, taking limits in the weak-operator topology we have that $P\pi(A)|_{\mathcal{H}_B} \subseteq \pi_B(B)$ and we may define $E : A \rightarrow B$ by

$$P\pi(a)|_{\mathcal{H}_B} = P\pi(E(a))|_{\mathcal{H}_B}.$$

Clearly $E(b) = b$ if $b \in B$ and $\|E(a)\| \leq \|a\| \forall a \in A$. Also, $\phi(E(a)) = \langle \pi_B(E(a))\xi, \xi \rangle = \langle P\pi(a)P\xi, \xi \rangle = \langle \pi(a)\xi, \xi \rangle = \phi(a)$, so $\phi \circ E = \phi$. The s.o.-continuity of E is clear. $\square$

Lemma 3.2 *Suppose (A, ϕ) is a C^* -algebra with faithful state. Suppose $B \subseteq A$ is a C^* -subalgebra with projection of norm 1, $E : A \rightarrow B$, such that $\phi \circ E = \phi$. Let $(\pi, \mathcal{H}, \xi) = \text{GNS}(A, \phi)$, let $\mathcal{H}_B = \overline{\{\hat{b} \mid b \in B\}}$ and let $P : \mathcal{H} \rightarrow \mathcal{H}_B$ be the orthogonal projection, with notation as in Lemma 3.1. Then*

$$(E(a))^\wedge = P\hat{a}, \tag{7}$$

$$P\pi(a)|_{\mathcal{H}_B} = P\pi(E(a))|_{\mathcal{H}_B} \quad \forall a \in A. \tag{8}$$

Proof The identity (7) follows because

$$\langle E(a)^\wedge, \hat{b} \rangle = \phi(b^* E(a)) = \phi(E(b^* a)) = \phi(b^* a) = \langle \hat{a}, \hat{b} \rangle = \langle \hat{a}, P\hat{b} \rangle = \langle P\hat{a}, \hat{b} \rangle \quad \forall b \in B$$

and then (8) holds because

$$P(\pi(a)\hat{b}) = P((ab)^\wedge) = E(ab)^\wedge = (E(a)b)^\wedge = E(a)\hat{b}.$$

$\square$

Proposition 3.3 *Suppose (A_ι, ϕ_ι) are von Neumann algebras with normal, faithful states, for ι in some index set I . Suppose $1 \in B_\iota \subseteq A_\iota$ are von Neumann subalgebras with projections of norm 1, $E_\iota : A_\iota \rightarrow B_\iota$, such that $\phi_\iota \circ E_\iota = \phi_\iota$. Let $(\mathcal{M}, \phi) = \bigstar_{\iota \in I} (A_\iota, \phi_\iota)$ and let $\mathcal{N} = W^*(\bigcup_{\iota \in I} B_\iota) \subseteq \mathcal{M}$. Then there is a projection of norm 1, $E : \mathcal{M} \rightarrow \mathcal{N}$ such that $\phi \circ E = \phi$ and $E(a) = E_\iota(a)$ whenever $a \in A_\iota$.*

Proof Let $(\pi_\iota, \mathcal{H}_\iota, \xi_\iota) = \text{GNS}(A_\iota, \phi_\iota)$, $(\pi, \mathcal{H}, \xi) = \text{GNS}(\mathcal{M}, \phi)$. From the construction of the free product we have that

$$\mathcal{H} = \mathbf{C}\xi \oplus \bigoplus_{\substack{n \geq 1 \\ \iota_1 \neq \iota_2 \neq \dots \neq \iota_n}} \overset{\circ}{\mathcal{H}}_{\iota_1} \otimes \dots \otimes \overset{\circ}{\mathcal{H}}_{\iota_n},$$

where $\overset{\circ}{\mathcal{H}}_\iota = \mathcal{H}_\iota \ominus \mathbf{C}\xi_\iota$. By Lemma 3.2 there is a projection $P_\iota : \mathcal{H}_\iota \rightarrow \mathcal{H}_{B_\iota}$ implementing the conditional expectation E_ι . If $\Omega = \Lambda^\circ((B_\iota^\circ)_{\iota \in I})$ then $\mathfrak{A} = \text{span}(\Omega \cup \{1\})$ is the span of words in the B_ι and is a s.o.-dense $*$ -subalgebra of $\mathcal{N}$. Thus it is clear that

$$\overline{\{\hat{a} \mid a \in \mathcal{N}\}} = \overline{\{\hat{a} \mid a \in \mathfrak{A}\}} = \mathcal{H}_{\mathcal{N}} = \mathbf{C}\xi \oplus \bigoplus_{\substack{n \geq 1 \\ \iota_1 \neq \iota_2 \neq \dots \neq \iota_n}} \overset{\circ}{\mathcal{H}}_{B_{\iota_1}} \otimes \dots \otimes \overset{\circ}{\mathcal{H}}_{B_{\iota_n}} \subseteq \mathcal{H}.$$

Let $P : \mathcal{H} \rightarrow \mathcal{H}_{\mathcal{N}}$ be the orthogonal projection. Then for a word $x = a_1 a_2 \dots a_n \in \Omega$, $a_j \in A_{\iota_j}$, $\iota_j \neq \iota_{j+1}$, we have that

$$P\pi(x)|_{\mathcal{H}_{\mathcal{N}}} = P\pi(E_{\iota_1}(a_1)E_{\iota_2}(a_2)\dots E_{\iota_n}(a_n))|_{\mathcal{H}_{\mathcal{N}}},$$

hence the same holds for all $x \in \Omega$. Thus, by Lemma 3.1, there is a conditional expectation E with the desired properties. $\square$

Corollary 3.4 *Suppose (A_ι, ϕ_ι) are von Neumann algebras with normal, faithful states, $(\iota = 1, 2)$, and let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$. Then there is a projection, E , of norm 1 from $\mathcal{M}$ onto A_1 such that $\phi_1 \circ E = \phi$. Moreover, E is continuous from the strong-operator topology on $\mathcal{M}$ (in its GNS representation corresponding to ϕ) to the strong-operator topology on A_1 (in its GNS representation corresponding to ϕ_1).*

4. Extremal almost periodic states.

Let us first recall a few definitions and theorems of Connes'.

Definition 4.1 (Connes [1974]) Let $(\mathcal{M}, \phi)$ consist of a von Neumann algebra, $\mathcal{M}$, with normal, faithful (n.f.) state or semi-finite weight, ϕ . Then ϕ is said to be *almost periodic* if the modular operator, Δ_ϕ , has pure point spectrum. If, moreover, $\mathcal{M}$ is full, one defines $Sd(\mathcal{M})$ to be the intersection of the point spectra of the modular operators Δ_ϕ as ϕ ranges over all n.f. almost periodic weights.

Almost periodic states behave well with respect to free products. We will be able to determine explicitly many free products that are taken with respect to almost periodic states. The following basic observation follows easily from Voiculescu's description of the conjugation operator (see Lemma 1.8 of Voiculescu [1985] or the discussion in §1 of Dykema [1994(b)]).

Proposition 4.2 *For each $i \in I$, given a von Neumann algebra, A_i , with n.f. almost periodic state, ϕ_i , whose modular operator has point spectrum Γ_i , consider the free product $(\mathcal{M}, \phi) = \ast_{i \in I} (A_i, \phi_i)$. Then ϕ is an almost periodic state and the point spectrum of its modular operator equals the subgroup of $\mathbf{R}_+^*$ generated by $\bigcup_{i \in I} \Gamma_i$.*

Remark 4.3 It follows from Arveson's and Connes' spectral theory (Arveson [1974], Connes [1973]) that whenever ϕ is an almost periodic n.f. state or weight on any von Neumann algebra $\mathcal{M}$ one has a decomposition

$$\mathcal{M} = \bigoplus_{\gamma \in \Gamma} \mathcal{M}_\phi(\gamma), \tag{9}$$

where Γ is the point spectrum of Δ_ϕ and where

$$\mathcal{M}_\phi(\gamma) = \{a \in \mathcal{M} \mid \phi(xa) = \gamma\phi(ax) \ \forall x \in \mathcal{M}\}.$$

(See §1 of Dykema [1995(b)].) One also has that each $\mathcal{M}_\phi(\gamma)$ is w.o.-closed, that $\mathcal{M}_\phi(\gamma_1)\mathcal{M}_\phi(\gamma_2) \subseteq \mathcal{M}_\phi(\gamma_1\gamma_2)$ and that $\mathcal{M}(\gamma)^* = \mathcal{M}(\gamma^{-1})$. Thus if $\gamma > 1$ and if $\mathcal{M}_\phi(\gamma)$ contains an isometry v , then $\phi(vv^*) = \gamma$ and $\mathcal{M}_\phi(\gamma) = \mathcal{M}_\phi v$, where $\mathcal{M}_\phi$ denotes the *centralizer* of ϕ , which is by definition $\mathcal{M}_\phi = \mathcal{M}_\phi(1)$.

Definition 4.4 Suppose ϕ is an almost periodic weight on a von Neumann algebra $\mathcal{M}$. Let Γ be the subgroup of $\mathbf{R}_+^*$ generated by the point spectrum of the modular operator Δ_ϕ . Suppose Γ' is a subgroup of Γ . Then define $\mathcal{M}_\phi(\Gamma')$ to be the weak closure of the linear span of $\bigcup_{\gamma \in \Gamma'} \mathcal{M}_\phi(\gamma)$.

Lemma 4.5 *In the situation of Definition 4.4, there is a projection of norm 1, E , from $\mathcal{M}$ onto $\mathcal{M}_\phi(\Gamma')$ such that $\phi \circ E = \phi$ and that is continuous from the strong-operator topology on $\mathcal{M}$ (in its GNS representation from ϕ) to the strong-operator topology on $\mathcal{M}_\phi(\Gamma')$ (in its GNS representation from ϕ). We have for $a \in \mathcal{M}_\phi(\gamma)$ that*

$$E(a) = \begin{cases} a & \text{if } \gamma \in \Gamma' \\ 0 & \text{otherwise.} \end{cases}$$

Proof Let $\mathcal{N}$ denote $\mathcal{M}_\phi(\Gamma')$. Let $(\pi, \mathcal{H}, \xi) = \text{GNS}(\mathcal{M}, \phi)$. Since $\text{span}(\bigcup_{\gamma \in \Gamma} \mathcal{M}_\phi(\gamma))$ is s.o.-dense in $\mathcal{M}$, we get an orthogonal decomposition of the Hilbert space, $\mathcal{H} = \bigoplus_{\gamma \in \Gamma} \mathcal{H}_\phi(\gamma)$, where $\mathcal{H}_\phi(\gamma) = \overline{\{\hat{a} \mid a \in \mathcal{M}_\phi(\gamma)\}}$, and one sees that $\mathcal{H}_\mathcal{N} = \bigoplus_{\gamma \in \Gamma'} \mathcal{H}_\phi(\gamma)$. If $P : \mathcal{H} \rightarrow \mathcal{H}_\mathcal{N}$ is the orthogonal projection then for $a \in \mathcal{M}_\phi(\gamma)$

$$P\pi(a)|_{\mathcal{H}_\mathcal{N}} = \begin{cases} P\pi(a)|_{\mathcal{H}_\mathcal{N}} & \text{if } \gamma \in \Gamma' \\ 0 & \text{otherwise.} \end{cases}$$

Hence Lemma 3.1 applies to give E with the desired properties. $\square$

Proposition 4.6 (Connes [1974]) *Suppose $\mathcal{M}$ is a full factor. Then $Sd(\mathcal{M})$ is a multiplicative subgroup of the positive reals, $\mathbf{R}_+^*$. If $Sd(\mathcal{M}) \neq \mathbf{R}_+^*$ and if ϕ is an almost periodic weight on $\mathcal{M}$, then (the point spectrum of Δ_ϕ) = $Sd(\mathcal{M})$ if and only if $\mathcal{M}_\phi$ is a factor.*

The following is in analogy with Connes' result.

Definition 4.7 An *extremal almost periodic state* (or weight) on a von Neumann algebra $\mathcal{M}$ is a normal, faithful, almost periodic state (or weight), ϕ , whose centralizer, $\mathcal{M}_\phi$, is a factor. We define $Sd(\phi)$ to be the point spectrum of the modular operator Δ_ϕ .

Lemma 4.8 *Suppose a von Neumann algebra $\mathcal{M}$ has an extremal almost periodic state ϕ . Then $\mathcal{M}$ is a factor and*

- (i) *is type II_1 if $Sd(\phi) = \{1\}$;*
- (ii) *is type III_λ if $Sd(\phi) = \{\lambda^n \mid n \in \mathbf{Z}\}$, $0 < \lambda < 1$;*
- (iii) *is type III_1 if $Sd(\phi)$ is dense in $\mathbf{R}_+^*$.*

Proof The center of $\mathcal{M}$ lies in the centralizer of ϕ , which is a factor, so $\mathcal{M}$ is a factor. The type classification (i), (ii), and (iii) of Connes' follows from Corollary 3.2.7. of Connes [1973]. $\square$

Lemma 4.9 *Suppose $\mathcal{M}$ is a factor with extremal almost periodic state ϕ . Then*

- (i) *for every $\gamma \in Sd(\phi)$, $\gamma > 1$, there is an isometry v_γ in the spectral subspace $\mathcal{M}_\phi(\gamma)$;*
- (ii) *$Sd(\phi)$ is a subgroup of the multiplicative group $\mathbf{R}_+^*$*

Proof Since γ is in the point spectrum of Δ_ϕ , one sees that $\mathcal{M}_\phi(\gamma)$ is nonempty. Taking the polar decomposition of any element of $\mathcal{M}_\phi(\gamma)$, one shows that $\mathcal{M}_\phi(\gamma)$ contains a partial isometry, v' . Then $\phi(v'v'^*) = \gamma\phi(v'^*v')$. Since $\mathcal{M}_\phi$ is a factor and $\gamma > 1$, there are partial isometries x_j and y_j in $\mathcal{M}_\phi$ such that $v = \sum_{j=1}^n x_j v' y_j \in \mathcal{M}_\phi(\gamma)$ is an isometry. This proves (i). Part (ii) now follows easily from (i) and the fact that $h_1 \mathcal{M}_\phi h_2 \neq \{0\}$ if h_1, h_2 are nonzero projections in $\mathcal{M}_\phi$.

Lemma 4.10

- (i) *Let $\mathcal{M}$ be a type III factor having an extremal almost periodic state ϕ and let $h \in \mathcal{M}_\phi$ be a self-adjoint projection. Then $\phi_h = \phi(h)^{-1}\phi|_{h\mathcal{M}h}$ is an extremal almost periodic state on $h\mathcal{M}h$ whose centralizer is equal to $h\mathcal{M}_\phi h$, and $Sd(\phi_h) = Sd(\phi)$.*
- (ii) *Conversely, if $\mathcal{M}$ is a von Neumann algebra with almost periodic state ϕ and if $h \in \mathcal{M}_\phi$ is a self-adjoint projection whose central support in $\mathcal{M}_\phi$ is 1 and such that $h\mathcal{M}_\phi h$ is a factor, then $\mathcal{M}$ is a factor and ϕ is extremal almost periodic.*

Proof Part (i) is clear from Lemma 4.9 and Remark 4.3 and part (ii) is tautologically true, because $\mathcal{M}_\phi$ must be a factor. $\square$

In the remainder of this section, we prove that an extremal almost periodic state, ϕ , is truly extremal, in the sense that if ψ is an almost periodic state that commutes with ϕ in the sense of Pedersen and Takesaki [1973], then the point spectrum of Δ_ψ contains the point spectrum of Δ_ϕ .

Given faithful, normal states, ϕ and ψ , on a von Neumann algebra $\mathcal{M}$, by Pedersen and Takesaki's Radon-Nikodym Theorem for von Neumann algebras, Pedersen and Takesaki [1973], ψ commutes with ϕ if and only if there is a self-adjoint, nonnegative operator, h , affiliated with the centralizer of ϕ , $\mathcal{M}_\phi$, such that $\psi = \phi(\cdot h)$.

Lemma 4.11 *Let ϕ be a faithful, extremal almost periodic, normal state on $\mathcal{M}$ and let ψ be a faithful, normal state on $\mathcal{M}$ commuting with ϕ , so $\psi = \phi(\cdot h)$ as described above. Then ψ is almost periodic if and only if h has purely atomic spectral measure, i.e. if and only if $h = \sum_{k \in I} t_k E_k$ for $t_k \in \mathbf{R}_+^*$ and $(E_k)_{k \in I}$ an orthogonal family of projections in $\mathcal{M}_\phi$ whose sum is 1.*

Proof Let $E(\cdot)$ be the spectral measure of h , so $h = \int_{\mathbf{R}_+} t E(dt)$. Let

$$\eta_\phi : \mathcal{M} \rightarrow L^2(\mathcal{M}, \phi)$$

$$\eta_\psi : \mathcal{M} \rightarrow L^2(\mathcal{M}, \psi)$$

be the usual embeddings, and denote the inner product in $L^2(\mathcal{M}, \phi)$ by $\langle \cdot, \cdot \rangle_\phi$ and in $L^2(\mathcal{M}, \psi)$ by $\langle \cdot, \cdot \rangle_\psi$. Let Γ_ϕ denote the point spectrum of Δ_ϕ and let

$$\mathfrak{m} = \bigcup_{\substack{0 < t_1 < t_2 < \infty \\ \gamma \in \Gamma_\phi}} E([t_1, t_2]) \mathcal{M}_\phi(\gamma) E([t_1, t_2]),$$

so $\mathfrak{m}$ is s.o.-dense in $\mathcal{M}$. Then for $x \in \mathfrak{m} \cap \mathcal{M}_\phi(\gamma)$ and $y \in \mathfrak{m}$,

$$\begin{aligned} \langle \eta_\psi(x), \eta_\psi(y) \rangle_\psi &= \psi(y^*x) = \phi(y^*xh) = \langle \eta_\phi(xh), \eta_\phi(y) \rangle_\phi \\ \langle \Delta_\psi \eta_\psi(x), \eta_\psi(y) \rangle_\psi &= \langle \eta_\psi(y^*), \eta_\psi(x^*) \rangle_\psi = \psi(xy^*) = \phi((hx)y^*) = \gamma\phi(y^*(hx)) = \\ &= \gamma\psi(y^*(h x h^{-1})) = \gamma\langle \eta_\psi(h x h^{-1}), \eta_\psi(y) \rangle_\psi. \end{aligned} \quad (10)$$

For $\gamma \in \Gamma_\phi$, let F_γ be the orthogonal projection from $L^2(\mathcal{M}, \psi)$ onto $\overline{\text{span}}\{\eta_\psi(x) \mid x \in \mathcal{M}_\gamma(\phi)\}$. Then $(F_\gamma)_{\gamma \in \Gamma_\phi}$ is an orthogonal family of projections whose sum is 1. Let $D_\phi = \sum_{\gamma \in \Gamma_\phi} \gamma F_\gamma$. Let L_h and $R_{h^{-1}}$ be the closures of the operators defined on $\mathfrak{m}$ by $L_h \eta_\psi(x) = \eta_\psi(hx)$ and $R_{h^{-1}} \eta_\psi(x) = \eta_\psi(xh^{-1})$. Clearly, $L_{E(\cdot)}$ is the spectral measure for L_h and $R_{E(\sigma(\cdot))}$ is the spectral measure for $R_{h^{-1}}$, where $\sigma(t) = t^{-1}$. Then D_ϕ , L_h and $R_{h^{-1}}$ are commuting, (see §VII.5 of Reed and Simon [1972]), nonnegative, self-adjoint operators on $L^2(\mathcal{M}, \psi)$. From (10) we see that $\Delta_\psi = D_\phi L_h R_{h^{-1}}$. Since $\mathcal{M}_\phi$ is a factor, and using the spectral decomposition of $\mathcal{M}$ with respect to ϕ , (see Remark 4.3), we see that $L^2(\mathcal{M}, \psi)$ has a basis of eigenvectors of Δ_ψ if and only if h has purely atomic spectral measure. $\square$

Proposition 4.12 *Let ϕ and ψ be faithful, almost periodic, normal states on $\mathcal{M}$ such that ψ commutes with ϕ in the sense of Pedersen and Takesaki [1973]. If ϕ is extremal almost periodic then the point spectrum of Δ_ψ contains the point spectrum of Δ_ϕ .*

Proof Let h , affiliated with $\mathcal{M}_\phi$, be such that $\psi = \phi(\cdot h)$. By Lemma 4.11 and its proof, h has purely atomic spectral measure and $\Delta_\psi = D_\phi L_h R_{h^{-1}}$. Let Γ_h be the point spectrum of h , so $h = \sum_{t \in \Gamma_h} t E_t$, where $(E_t)_{t \in \Gamma_h}$ is an orthogonal family of projections in $\mathcal{M}_\phi$ summing to 1. Using the spectral decomposition (see Remark 4.3) of $\mathcal{M}$ with respect to ϕ and the factoriality of $\mathcal{M}_\phi$, we see that $E_{t_1} F_\gamma E_{t_2}^{-1} \neq 0$ for every $\gamma \in \Gamma_\phi$ and every $t_1, t_2 \in \Gamma_h$, (where F_γ is as in the proof of 4.11). Thus the point spectrum of Δ_ψ is equal to $\{\gamma t_1 t_2^{-1} \mid \gamma \in \Gamma_\phi, t_1, t_2 \in \Gamma_h\} \supseteq \Gamma_\phi$. $\square$

5. Some technical lemmas.

The free etymology proof of Theorem 1.2 of Dykema [1993(b)], which was stated for the finite, separable case, is valid also more generally. We restate it for convenience in two steps, first the more often used version, then its generalization.

Proposition 5.1 *Suppose (A_ι, ϕ_ι) is a von Neumann algebra with normal, faithful state ($\iota = 1, 2$) and let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$. Suppose $h \in A_1$ is a central projection. Let $B_1 = \mathbf{C}h + (1-h)A_1$ and $\mathcal{N} = W^*(B_1 \cup A_2)$. Then $h\mathcal{M}h = W^*(h\mathcal{N}h \cup A_1h)$ and $h\mathcal{N}h$ and A_1h are free in $(h\mathcal{M}h, \phi(h)^{-1}\phi|_{h\mathcal{M}h})$. Moreover, the central support, $\mathcal{C}_\mathcal{M}(h)$, of h in $\mathcal{M}$ equals the central support, $\mathcal{C}_\mathcal{N}(h)$, of h in $\mathcal{N}$.*

Proposition 5.2 *Suppose (A_ι, ϕ_ι) is a von Neumann algebra with normal, faithful state $(\iota = 1, 2)$ and let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$. Suppose $A_1 = D_1 \otimes M_n(\mathbf{C}) \oplus D_2$ and let*

$$\begin{aligned} A_1 &= D_1 \otimes M_n(\mathbf{C}) \oplus D_2 \\ &\cup \\ B_1 &= 1 \otimes M_n(\mathbf{C}) \oplus D_2 \end{aligned}$$

and $\mathcal{N} = W^*(B_1 \cup A_2)$. Let $h \in 1 \otimes M_n(\mathbf{C}) \oplus 0 \subseteq B_1$ be a minimal projection. Then $h\mathcal{M}h = W^*(h\mathcal{N}h \cup (D_1 \otimes h))$ and $h\mathcal{N}h$ and $D_1 \otimes h$ are free in $(h\mathcal{M}h, \phi(h)^{-1}\phi|_{h\mathcal{M}h})$.

Lemma 5.3 *Suppose (A_ι, ϕ_ι) is a von Neumann algebra with normal, faithful state $(\iota = 1, 2)$ and let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$. Suppose $h_\iota \in A_\iota$ is a central projection and $\phi_1(h_1) = \phi_2(h_2)$. Let $B_\iota = \mathbf{C}h_\iota + (1 - h_\iota)A_\iota$ and $\mathcal{N} = W^*(B_1 \cup B_2)$. There is a unitary $w \in W^*(1, h_1, h_2)$ such that $h_2w = wh_1$. Then $h_1\mathcal{M}h_1 = W^*(h_1\mathcal{N}h_1 \cup A_1h_1 \cup w^*A_2h_2w)$ and $(h_1\mathcal{N}h_1, A_1h_1, w^*A_2h_2w)$ is free in $(h_1\mathcal{M}h_1, \phi(h_1)^{-1}\phi|_{h_1\mathcal{M}h_1})$. Moreover, the central supports satisfy $\mathcal{C}_{\mathcal{M}}(h_1) = \mathcal{C}_{\mathcal{N}}(h_1)$.*

Proof That the unitary w exists is easily seen from Theorem 1.1 of Dykema [1993(b)]. Let $\mathcal{N}_1 = W^*(B_1 \cup A_2)$. Then by Proposition 5.1, $h_2\mathcal{N}_1h_2 = W^*(h_2\mathcal{N}h_2 \cup h_2A_2)$, $h_2\mathcal{N}h_2$ and h_2A_2 are free in $h_2\mathcal{N}_1h_2$ and $\mathcal{C}_{\mathcal{N}_1}(h_2) = \mathcal{C}_{\mathcal{N}}(h_2)$. Thus $h_1\mathcal{N}_1h_1 = w^*h_2\mathcal{N}_1h_2w = W^*(h_1\mathcal{N}h_1 \cup w^*h_2A_2w)$, $h_1\mathcal{N}h_1$ and $w^*h_2A_2w$ are free in $h_1\mathcal{N}_1h_1$ and $\mathcal{C}_{\mathcal{N}_1}(h_1) = \mathcal{C}_{\mathcal{N}}(h_1)$. Applying again Proposition 5.1, we have $h_1\mathcal{M}h_1 = W^*(h_1\mathcal{N}_1h_1 \cup h_1A_1)$, $h_1\mathcal{N}_1h_1$ and h_1A_1 are free in $h_1\mathcal{M}h_1$ and $\mathcal{C}_{\mathcal{M}}(h_1) = \mathcal{C}_{\mathcal{N}_1}(h_1)$. Now apply Proposition 2.5.5 of Voiculescu, Dykema and Nica [1992]. $\square$

The previous two results were about what happens when one adds central summands in a free product situation (i.e. going from B_ι to A_ι) and making the center of A_ι larger (or keeping it the same). The next result is about what happens when adding a partial isometry and making the center of A_ι smaller.

Lemma 5.4 *Suppose (A_ι, ϕ_ι) is a von Neumann algebra with normal, faithful state $(\iota = 1, 2)$ and let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$. Suppose $v \in A_1$ is a partial isometry such that $vv^* = k$, $v^*v = h \leq 1 - k$, with $h, k \in (A_1)_{\phi_1}$ and*

$$\forall a \in A_1 \quad \phi_1(va) = 0 \Leftrightarrow \phi_1(av) = 0.$$

Set B_1 to be either

- (i) $B_1 = \mathbf{C}k + (1 - k)A_1(1 - k) \subseteq A_1$,
- (ii) or, only if k is minimal in A_1 , $B_1 = \tilde{k}A_1\tilde{k} + \tilde{h}A_1\tilde{h}$ where $\tilde{k}, \tilde{h} \in (A_1)_{\phi_1}$ are projections, $h \leq \tilde{h}$, $k \leq \tilde{k}$ and $\tilde{h} + \tilde{k} = 1$.

Important properties of B_1 will be that k is minimal in B_1 and that $kB_1h = \{0\}$. Let $\mathcal{N} = W^*(B_1 \cup A_2)$, let $\mathcal{N}_\phi$ denote the centralizer of $\phi|_{\mathcal{N}}$ and suppose there is $x \in \mathcal{N}$ such that $x^*x = h$, $xx^* = k$ and

$$\forall a \in \mathcal{N} \quad \phi(xa) = 0 \Leftrightarrow \phi(ax) = 0.$$

Then x^*v is a Haar unitary and $(h\mathcal{N}h, \{x^*v\})$ is $*$ -free in $(h\mathcal{M}h, \phi(h)^{-1}\phi|_{h\mathcal{M}h})$, and

$$h\mathcal{M}h = W^*(h\mathcal{N}h \cup \{x^*v\}). \tag{11}$$

Moreover, the central supports satisfy $\mathcal{C}_{\mathcal{M}}(h) = \mathcal{C}_{\mathcal{N}}(h + k)$.

Proof (11) is clear by the usual algebraic technique of bringing back generators with partial isometries, cf. Lemma 1 of Rădulescu [1992(a)]. We will show simultaneously that x^*v is a Haar unitary and that $(h\mathcal{N}h, \{x^*v\})$ is $*$ -free by showing that $\phi(y) = 0$ for every alternating product y in $(h\mathcal{N}h)^\circ$ and $\{(x^*v)^n \mid n \in \mathbf{N}\} \cup \{(v^*x)^n \mid n \in \mathbf{N}\}$. Regrouping by sticking x and x^* to letters from $(h\mathcal{N}h)^\circ$ whenever possible, we see that y is equal to an alternating product in $\{v, v^*\}$ and

$$(h\mathcal{N}h)^\circ \cup (h\mathcal{N}h)x^* \cup x(h\mathcal{N}h) \cup x(h\mathcal{N}h)^\circ x^*,$$

where

every letter from $(h\mathcal{N}h)^\circ$ is (unless it is the first letter of y) preceeded by v
and is (unless it is the last letter of y) followed by v^* ;

every letter from $(h\mathcal{N}h)x^*$ is (unless it is the first letter of y) preceeded by v
and is followed by v ;

every letter from $x(h\mathcal{N}h)$ is preceeded by v^*
and is (unless it is the last letter of y) followed by v^* ;

every letter from $x(h\mathcal{N}h)^\circ x^*$ is preceeded by v^*
and is followed by v .

Using Kaplansky's density theorem, we see that every element of $\mathcal{N}$ is the s.o.-limit of a bounded sequence in $\text{span}(\Omega \cup \{1\})$ where $\Omega = \Lambda^\circ(B_1^\circ, A_2^\circ)$. Now using the ϕ -preserving conditional expectation from $\mathcal{N}$ onto B_1 (which is gotten from Proposition 3.4), and using the fact that k is minimal in B_1 and k and h are perpendicular projections belonging to the centralizer of ϕ , we have that every element of

$(h\mathcal{N}h)^\circ$ is the s.o.-limit of a bounded sequence in $\text{span } \Omega_0$

$x(h\mathcal{N}h)$ is the s.o.-limit of a bounded sequence in $\text{span } \Omega_r$

$(h\mathcal{N}h)x^*$ is the s.o.-limit of a bounded sequence in $\text{span } \Omega_l$

$x(h\mathcal{N}h)^\circ x^*$ is the s.o.-limit of a bounded sequence in $\text{span } \Omega_{l,r}$

where

$$\Omega_0 = (\Omega \setminus B_1^\circ) \cup (hB_1h)^\circ$$

Ω_r is the set of elements of Ω whose right-most letter is from A_2°

Ω_l is the set of elements of Ω whose left-most letter is from A_2°

$$\Omega_{l,r} = \Omega_l \cap \Omega_r.$$

Hence it will suffice to show that $\phi(z) = 0$ whenever z is an alternating product in $\{v, v^*\}$ and $\Omega_0 \cup \Omega_l \cup \Omega_r \cup \Omega_{l,r}$, such that every letter that is v is preceded by a letter from $\Omega_r \cup \Omega_{l,r}$ and every letter that is v^* is followed by a letter from $\Omega_l \cup \Omega_{l,r}$. Regrouping by sticking each v and v^* to a letter from B_1° whenever possible, and using that

$v(hB_1h)^\circ v^* \subseteq A_1^\circ$ and $vB_1 \subseteq A_1^\circ$, we see that z equals an alternating product in A_1° and A_2° , hence by freeness, $\phi(z) = 0$.

We have left to prove only that $\mathcal{C}_\mathcal{M}(h) = \mathcal{C}_\mathcal{N}(h+k)$. Using $\mathcal{C}_\mathcal{A}(p) = \bigwedge_{x \in \mathcal{A}} \text{left support}(xp)$, and noting $vh = v$ has left support equal to k , we have that $\mathcal{C}_\mathcal{M}(h) = \mathcal{C}_\mathcal{M}(h+k) \geq \mathcal{C}_\mathcal{N}(h+k)$. But since $\mathcal{C}_\mathcal{N}(h+k)$ commutes with B_1 , with A_2 and with v , we have that $\mathcal{C}_\mathcal{N}(h+k)$ is in the center of $\mathcal{M}$, hence $\mathcal{C}_\mathcal{M}(h+k) \leq \mathcal{C}_\mathcal{N}(h+k)$. $\square$

Now we prove that (in certain circumstances) if A_1 is freely complemented in $\mathcal{M}$ then hA_1h is freely subcomplemented in $h\mathcal{M}h$.

Proposition 5.5 *Let A_ι be a von Neumann algebra and ϕ_ι a normal, faithful state on A_ι , ($\iota = 1, 2$). Let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$. Suppose $(A_1)_{\phi_1}$ is a factor and $(A_2)_{\phi_2}$ has no minimal projections. If $1 \neq h \in (A_1)_{\phi_1}$ is a self-adjoint projection then hA_1h is freely subcomplemented in $(h\mathcal{M}h, \phi(h)^{-1}\phi|_{h\mathcal{M}h})$ by $(B_\iota)_{0 \leq \iota \leq n}$, some $n \geq 1$, where B_0 is isomorphic to the cut-down of the free product of A_2 and a finite dimensional abelian algebra and where $B_\iota \cong L(\mathbf{Z}) \ \forall \iota \geq 1$.*

Proof There are $n \geq 1$ and partial isometries $v_0, v_1, \dots, v_n \in (A_1)_{\phi_1}$ such that $v_j v_j^* \leq h \ \forall j$ and $\sum_{j=0}^n v_j^* v_j = 1$. We may assume $v_0 = h$. Let

$$\begin{aligned} D_{-1} &= \mathbf{C}h + \mathbf{C}v_1^* v_1 + \dots + \mathbf{C}v_n^* v_n \subseteq (A_1)_{\phi_1} \\ D_0 &= D_{-1} + hA_1h \subseteq A_1 \\ D_j &= W^*(D_{j-1} \cup \{v_j\}) \subseteq A_1 \text{ for } 1 \leq j \leq n. \end{aligned}$$

For $-1 \leq j \leq n$ let $\mathcal{P}_j = W^*(D_j \cup A_2)$ so $(\mathcal{P}_j, \phi|_{\mathcal{P}_j}) \cong (D_j, \phi_1|_{D_j}) * (A_2, \phi_2)$. Clearly $D_n = A_1$ and $\mathcal{P}_n = \mathcal{M}$. By Proposition 5.1, hA_1h is freely complemented in $h\mathcal{P}_0h$ by $h\mathcal{P}_{-1}h$. From Dykema [1993(b)] we see that there are partial isometries x_j in the centralizer of ϕ restricted to $W^*(D_{-1} \cup (A_2)_\phi)$ such that $x_j^* x_j = v_j^* v_j$ and $x_j x_j^* = v_j v_j^*$, so by Lemma 5.4 $h\mathcal{P}_{j-1}h$ is freely subcomplemented in $h\mathcal{P}_j h$ by an algebra isomorphic to $L(\mathbf{Z})$ with associated projection equal to $v_j v_j^* \in (A_1)_{\phi_1} \ \forall 1 \leq j \leq n$. Thus using Corollary 1.7 and the fact that $h(A_1)_{\phi_1} h$ is a factor the proposition is proved. $\square$

Now we prove that (in certain circumstances) if A is a freely subcomplemented von Neumann subalgebra of $\mathcal{M}$ then hAh is freely subcomplemented in $h\mathcal{M}h$.

Proposition 5.6 *Let $\mathcal{M}$ be a von Neumann algebra with normal, faithful state ϕ and $1 \in A \in \mathcal{M}$ a von Neumann subalgebra. Let A_ϕ denote the centralizer of $\phi|_A$ and suppose A_ϕ is a II_1 -factor. Suppose also that A is freely subcomplemented in $\mathcal{M}$ by $(B_\iota)_{\iota \in I}$ where for each $\iota \in I$ the centralizer of $\phi|_{B_\iota}$ has no minimal projections. If $h \in A_\phi$ is a self-adjoint projection then hAh is freely subcomplemented in $h\mathcal{M}h$ by algebras isomorphic to the B_ι , to cut-downs of free products of the B_ι with finite dimensional abelian algebras and to $L(\mathbf{Z})$.*

Proof Denote the associated projections of $(B_\iota)_{\iota \in I}$ by $(h_\iota)_{\iota \in I}$. Let us first do the case when $|I| = 1$, say $I = \{1\}$. Since A_ϕ as a factor we may assume either $h_1 \leq h$ or $h \leq h_1$. If $h_1 \leq h$ then clearly hAh is freely subcomplemented in $h\mathcal{M}h$ by B_1 . If $h \leq h_1$ then $h_1 A h_1$ and B_1 freely generate $h_1 \mathcal{M} h_1$ so applying Proposition 5.5 we see that hAh is

freely complemented in $h\mathcal{M}h$ by a cut-down of $(B_1$ free product a finite dimensional abelian algebra) and copies of $L(\mathbf{Z})$, so the case $|I| = 1$ is proved. Moreover in the above case, since A_ϕ is a II_1 -factor we may choose the associated projections for the free subcomplement of hAh in $h\mathcal{M}h$ to be in any given II_1 -subfactor of hAh . This allows us to conclude the general case from Corollary 1.8. $\square$

6. Free products of factors with respect to extremal almost periodic states.

Lemma 6.1 *Suppose ϕ_ι is an almost periodic state on a von Neumann algebra A_ι , ($\iota = 1, 2$). Let $\Gamma_{\iota,0}$ be the point spectrum of ϕ_ι and let Γ_ι be the subgroup of $\mathbf{R}_+^*$ generated by $\Gamma_{\iota,0}$. Let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$. Then by Proposition 4.2, ϕ is almost periodic and its point spectrum, which we denote Γ , is equal to the subgroup of $\mathbf{R}_+^*$ generated by $\Gamma_1 \cup \Gamma_2$. Suppose that $\lambda \in \Gamma_{1,0}$, $0 < \lambda < 1$, that there is an isometry $v \in (A_1)_{\phi_1}(\lambda^{-1})$ and that $\lambda^{\mathbf{Z}} \stackrel{\text{def}}{=} \{\lambda^n \mid n \in \mathbf{Z}\}$ is complemented in Γ_1 , say¹ $\Gamma_1 = H \oplus \lambda^{\mathbf{Z}}$. Let $B_1 = (A_1)_{\phi_1}(H)$. Note by Remark 4.3 that $\mathcal{M}$ is generated by B_1 and A_2 together with v . Let*

$$\begin{aligned} \mathcal{N}_{[0,0]} &= W^*(B_1 \cup A_2), \\ \mathcal{N}_{[-n,0]} &= W^*(v^* \mathcal{N}_{[-n+1,0]} v \cup \mathcal{N}_{[0,0]}), \quad n \geq 1, \\ \mathcal{N}_{(-\infty,0]} &= W^*(\bigcup_{n \geq 0} \mathcal{N}_{[-n,0]}), \\ \mathcal{N}_{(-\infty,n]} &= W^*(\mathcal{N}_{(-\infty,n-1]} \cup v^n \mathcal{N}_{[0,0]} (v^*)^n), \quad n \geq 1, \\ \mathcal{N}_{(-\infty,\infty)} &= W^*(\bigcup_{n \geq 0} \mathcal{N}_{(-\infty,n]}). \end{aligned} \tag{13}$$

Then

- (i) $v^* \mathcal{N}_{[-n+1,0]} v$ and $\mathcal{N}_{[0,0]}$ are free in $(\mathcal{M}, \phi)$ with amalgamation over B_1 , $\forall n \geq 1$;
- (ii) $h_n \mathcal{N}_{(-\infty,n-1]} h_n$ and $v^n \mathcal{N}_{[0,0]} (v^*)^n$ are free in $(h_n \mathcal{M} h_n, \lambda^{-n} \phi|_{h_n \mathcal{M} h_n})$ with amalgamation over $h_n B_1 h_n$, $\forall n \geq 1$, where $h_n = v^n (v^*)^n \in \mathcal{M}_\phi$;
- (iii) if $\lambda^{\mathbf{Z}} \cap \Gamma' = \{1\}$, where Γ' is the subgroup of Γ generated by H together with Γ_2 , then $\mathcal{N}_{(-\infty,\infty)} = \mathcal{M}_\phi(\Gamma')$.

Proof We have by Remark 4.3 that $\text{span}(B_1 \cup \bigcup_{n \in \mathbf{N}} (B_1 v^n \cup (v^*)^n B_1))$ is a s.o.-dense $*$ -subalgebra of A_1 , hence, letting

$$\Omega = \Lambda^\circ(B_1^\circ \cup \bigcup_{n \in \mathbf{N}} (B_1 v^n \cup (v^*)^n B_1), A_2^\circ),$$

we have that $\text{span}(\Omega \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{M}$. Consider a word $x = a_1 a_2 \cdots a_n$ in Ω , written as an alternating product of the specified form, i.e. each $a_j \in B_1^\circ \cup \bigcup_{n \in \mathbf{N}} (B_1 v^n \cup (v^*)^n B_1) \subseteq A_1^\circ$ or $a_j \in A_2^\circ$, with $a_j \in A_1^\circ \iff a_{j+1} \in A_2^\circ$. We will want to keep track of the occurrences of v and v^* , hence let $l(x) = n$ be the length of x and for each $1 \leq i, j \leq n$ let

$$\begin{aligned} s(a_i) &= \begin{cases} 0 & \text{if } a_i \in B_1^\circ \cup A_2^\circ \\ k & \text{if } a_i \in B_1 v^k \\ -k & \text{if } a_i \in (v^*)^k B_1, \end{cases} \\ t_j(x) &= \sum_{i=1}^j s(a_i). \end{aligned}$$

¹We will mostly use additive notation for direct sums, complements and cosets of subgroups of $\mathbf{R}_+^*$, even though the operation in $\mathbf{R}_+^*$ is multiplication.

If $0 \in I \subseteq \mathbf{Z}$ let Θ_I be the set of alternating products $x \in \Omega$ such that $t_{l(x)}(x) = 0$ and $t_j(x) \in I \forall 1 \leq j \leq l(x)$.

Claim 6.1a *If I is of the form $[-n, 0]$, $(-\infty, n]$, $n \in \{0\} \cup \mathbf{N}$, or $(-\infty, \infty)$ then $\text{span}(\Theta_I \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{N}_I$.*

Proof First, show that $\text{span}(\Theta_I \cup \{1\})$ is a $*$ -algebra. We need only show that multiplication is closed. We show by induction on $n + m$ that if we have elements of Ω , $x = a_1 \cdots a_n$ (or $x = 1$, in which case we take $n = 0$) and $y = b_1 \cdots b_m$ (or $y = 1$, in which case $m = 0$), such that $t_j(x) \in I$, $t_{l(x)}(x) + t_j(y) \in I \forall j$ and $t_n(x) + t_m(y) = 0$, then $xy \in \text{span}(\Theta_I \cup \{1\})$. When $n + m = 0$ or 1 , then either $x = 1$ or $y = 1$, so $xy \in \Theta_I \cup \{1\}$. If $a_n \in A_{\iota_1}^\circ$, $b_1 \in A_{\iota_2}^\circ$, $\iota_1 \neq \iota_2$, then $xy = a_1 \cdots a_n b_1 \cdots b_m$ is already in reduced form, so xy belongs to Θ_I . If $a_n \in A_{\iota}^\circ$ and $b_1 \in A_{\iota}^\circ$, let $c = a_n b_1$. If $\iota = 1$ and $s(a_n) \neq -s(b_1)$ then $a_n b_1 \in (v^*)^k B_1$ or $b_1 v^k$, some $k > 0$, so $xy = a_1 \cdots a_{n-1} c b_2 \cdots b_m$ is in reduced form and

$$t_j(xy) = \begin{cases} t_j(x) & \text{if } 1 \leq j \leq n-1 \\ t_{l(x)}(x) + t_{j-n+1}(y) & \text{if } n \leq j \leq n+m-1, \end{cases}$$

hence xy is an element of Θ_I . If $\iota = 1$ and $s(a_n) = -s(b_1)$ then $c \in B_1$, or if $\iota = 2$ then $c \in A_2$ and $c = c^\circ + \phi(c)1$ where $\phi(c^\circ) = 0$, so

$$xy = a_1 \cdots a_{n-1} c^\circ b_2 \cdots b_m + \phi(c) a_1 \cdots a_{n-1} b_2 \cdots b_m,$$

and, by the same argument as above, the first term is an element of Θ_I while the second term is of the correct form to apply the inductive hypothesis. Hence we have shown that $\text{span}(\Theta_I \cup \{1\})$ is a $*$ -algebra.

Now we show by induction on $n \geq 0$ that

$$\mathcal{N}_{[-n,0]} = \overline{\text{span}}(\Theta_{[-n,0]} \cup \{1\}). \quad (14)$$

First, note that $\mathcal{N}_{[-n,0]} \supseteq \mathcal{N}_{[-n+1,0]} \forall n \geq 1$, (which is readily shown from (13) by induction on n). For $n = 0$, (14) is clearly true. Suppose $n \geq 1$. By inductive hypothesis, we have $v^* \mathcal{N}_{[-n+1,0]} v \subseteq \overline{\text{span}}(v^* \Theta_{[-n+1,0]} v \cup \{1\}) \subseteq \overline{\text{span}}(\Theta_{[-n,0]} \cup \{1\})$, and $\mathcal{N}_{[0,0]} \subseteq \overline{\text{span}}(\Theta_{[0,0]} \cup \{1\})$, so the inclusion $\subseteq$ in (14) is true. For the reverse inclusion, suppose $x = a_1 \cdots a_m \in \Theta_{[-n,0]}$ is a word of the specified form. We show by induction on m that $x \in \mathcal{N}_{[-n,0]}$. If $t_j = 0 \forall 1 \leq j \leq m$ then $x \in \mathcal{N}_{[0,0]}$. Otherwise, let j_1 be least such that $t_{j_1}(x) < 0$ and j_2 least such that $j_2 > j_1$ and $t_{j_2}(x) = 0$. If $j_1 > 1$ then $a_1 \cdots a_{j_1-1} \in \mathcal{N}_{[0,0]}$. Now $a_{j_1} = (v^*)^{k_1} b_1$ and $a_{j_2} = b_2 v^{k_2}$, some $b_1, b_2 \in B_1$, $0 < k_1, k_2 \leq n$. Thus $(v^*)^{k_1-1} b_1 a_{j_1+1} \cdots a_{j_2-1} b_2 v^{k_2-1} \in \Theta_{[-n+1,0]}$, so (by inductive hypothesis for the induction on n), $a_{j_1} \cdots a_{j_2} \in v^* \mathcal{N}_{[-n+1,0]} v$. Finally, $a_{j_2+1} \cdots a_m \in \Theta_{[-n,0]}$, and is of shorter length, hence lies in $\mathcal{N}_{[-n,0]}$, (by inductive hypothesis for the induction on m). Thus $a_1 \cdots a_m \in \mathcal{N}_{[-n,0]}$, and (14) is true.

Let us now prove by induction on $n \geq 0$ that

$$\mathcal{N}_{(-\infty, n]} = \overline{\text{span}}(\Theta_{(-\infty, n]} \cup \{1\}). \quad (15)$$

For $n = 0$ we have it by taking the inductive limit of (14). The inclusion $\subseteq$ in (15) is true because (using inductive hypothesis) $\mathcal{N}_{(-\infty, n-1]} \subseteq \overline{\text{span}}(\Theta_{(-\infty, n-1]} \cup \{1\})$ and $v^n \Theta_{[0,0]} (v^*)^n \subseteq \Theta_{(-\infty, n]}$, while every element of $\mathcal{N}_{[0,0]}$ is the s.o.-limit of a bounded sequence in $\text{span}(\Theta_{[0,0]} \cup \{1\})$. To show $\supseteq$, consider $x = a_1 \cdots a_m \in \Theta_{(-\infty, n]}$, a word of the specified form, and show by induction on m that $x \in \mathcal{N}_{(-\infty, n]}$. If $t_j(x) < n \forall j$, then by inductive hypothesis (for the induction on n), $x \in \mathcal{N}_{(-\infty, n-1]} \subseteq \mathcal{N}_{(-\infty, n]}$. Otherwise let j_1 be least such that $t_{j_1}(x) = n$ and j_2 least

such that $j_2 > j_1$ and $t_{j_2}(x) < n$. Thus $a_{j_1} = b_1 v^{k_1}$ and $a_{j_2} = (v^*)^{k_2} b_2$ for some $b_1, b_2 \in B_1$ and $k_1, k_2 > 0$. Then $x = a_1 \cdots a_{j_1} (v^*)^n v^n a_{j_1+1} \cdots a_{j_2-1} (v^*)^n v^n a_{j_2} \cdots a_m$. But $a_1 \cdots a_{j_1} (v^*)^n \in \text{span} \Theta_{(-\infty, n-1]} \subseteq \mathcal{N}_{(-\infty, n-1]}$, $v^n a_{j_1+1} \cdots a_{j_2-1} (v^*)^n \in v^n (\Theta_{[0,0]} \cup \{1\}) (v^*)^n \subseteq v^n \mathcal{N}_{[0,0]} (v^*)^n$ and $v^n a_{j_2} \cdots a_m \in \text{span} \Theta_{(-\infty, n]}$ is a sum of one or two words of shorter length, so by inductive hypothesis, is in $\mathcal{N}_{(-\infty, n]}$. This proves (15).

Finally,

$$\mathcal{N}_{(-\infty, \infty)} = \overline{\text{span}}(\Theta_{(-\infty, \infty)} \cup \{1\})$$

is obtained by taking the inductive limit of (15). Hence Claim 6.1a is proved.

We continue with the proof of Lemma 6.1. Showing (i) is the same as showing that $\phi(z) = 0$ whenever z is an alternating product in $v^* \mathcal{N}_{[-n+1,0]} v \ominus B_1$ and $\mathcal{N}_{[0,0]} \ominus B_1$. If $x \in \mathcal{N}_{[0,0]}$ then (by Claim 6.1a, Kaplansky's density theorem and separability) x is the strong-operator limit of a bounded sequence, $(x_n)_{n=1}^\infty$ in $\text{span}(\Theta_{[0,0]} \cup \{1\})$. Combining Corollary 3.4 and Lemma 4.5, there is a ϕ -preserving conditional expectation, E_{B_1} from $\mathcal{M}$ onto B_1 . Hence if also $x \perp B_1$, then $E_{B_1}(x) = 0$ and thus $x_n - E_{B_1}(x_n)$ is a bounded sequence converging in s.o.-topology to x . So x is the s.o.-limit of a bounded sequence in $\text{span}(\Theta_{[0,0]} \setminus B_1^\circ)$. Doing the same for the other, note that $v^* \mathcal{N}_{[-n+1,0]} v \ominus B_1 = v^*(h_1 \mathcal{N}_{[-n+1,0]} h_1 \ominus h_1 B_1 h_1) v$. One similarly shows that if $x \in \mathcal{N}_{[-n+1,0]} \ominus B_1$ then x is the s.o.-limit of a bounded sequence in $\text{span}(\Theta_{[-n+1,0]} \setminus B_1^\circ)$. Hence, to show that $\phi(z) = 0$ it will suffice to show that $\phi(y) = 0$ whenever y is an alternating product in $\Theta_{[0,0]} \setminus B_1^\circ$ and $v^*(\Theta_{[-n+1,0]} \setminus B_1^\circ) v$. If we "erase parenthesis" in y and multiply elements from A_1 together whenever possible, because $B_1 v B_1 \subseteq A_1^\circ$ we see that y is equal to an alternating product in A_1° and A_2° , hence $\phi(y) = 0$ by freeness, and (i) is proved.

Part (ii) is proved similarly. It suffices to show that $\phi(y) = 0$ whenever y is an alternating product in $\Theta_{(-\infty, n-1]} \setminus B_1^\circ$ and $v^n(\Theta_{[0,0]} \setminus B_1^\circ)(v^*)^n$. Since every element of $\Theta_{(-\infty, n-1]} \setminus B_1^\circ$ begins with an element of either A_2° , $(v^*)^j B_1$, ($j \geq 1$), B_1° or $B_1 v^k$, ($1 \leq k \leq n-1$), and ends with the adjoint of one of these, by regrouping and multiplying neighbors we see that y is equal to an alternating product in A_1° and A_2° . Hence $\phi(y) = 0$ and (ii) is proved.

Let us now show (iii). We assume $\lambda^{\mathbf{Z}} \cap \Gamma' = \{1\}$. In light of Claim 6.1a, we have $\mathcal{N}_{(-\infty, \infty)} = \overline{\text{span}}(\Theta_{\mathbf{Z}} \cup \{1\})$. Note that $\Theta_{\mathbf{Z}}$ equals the set of words x belonging to Ω such that $t_{l(x)}(x) = 0$. Let E denote the projection from $\mathcal{M}$ onto $\mathcal{M}_\phi(\Gamma')$, obtained from Lemma 4.5. If $x \in \Omega$, then, writing each element of A_2° as the s.o.-limit of a bounded sequence in $\text{span} \bigcup_{\gamma \in Sd(\phi_2)} (A_2)_{\phi_2}(\gamma)$ and each element of B_1 as the s.o.-limit of a bounded sequence in $\text{span} \bigcup_{\gamma \in H} (A_1)_{\phi_1}(\gamma)$, we see that $x \in \overline{\text{span}} \bigcup_{\gamma \in \lambda^{k(x)} + \Gamma'} \mathcal{M}_\phi(\gamma)$, where $k(x) = -t_{l(x)}(x)$. So $t_{l(x)}(x) = 0$ implies $x \in \mathcal{M}_\phi(\Gamma')$, and $t_{l(x)}(x) \neq 0$ implies $x \perp \mathcal{M}_\phi(\Gamma')$ and hence $E(x) = 0$. Thus we easily see $\mathcal{M}_\phi(\Gamma') \supseteq \overline{\text{span}} \Theta_{\mathbf{Z}}$. For the reverse inclusion, let $y \in \mathcal{M}_\phi(\Gamma')$. Since $\text{span}(\Omega \cup \{1\})$ is s.o.-dense in $\mathcal{M}$, by the Kaplansky density theorem and separability there is a bounded sequence $(y_n)_{n \geq 1}$ converging to y in s.o.-topology and such that $y_n \in \text{span}(\Omega \cup \{1\}) \forall n$. Then, since $E(y) = y$, we have that $(E(y_n))_{n \geq 1}$ is a bounded sequence converging strongly to y , and $E(y_n) \in \text{span}(\Theta_{\mathbf{Z}} \cup \{1\}) \forall n$. Thus (iii) is proved. □

Lemma 6.2 *Let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$, Γ_ι , $\Gamma_{\iota,0}$ and Γ be as in Lemma 6.1. Suppose $\lambda \in \Gamma_{1,0}$, $0 < \lambda < 1$, there is an isometry $v \in (A_1)_{\phi_1}(\lambda^{-1})$ and there is a subgroup $H \subseteq \Gamma_1$ that together with λ generates Γ_1 . Suppose*

$\lambda^{\mathbf{Z}} \cap H = \lambda^{N\mathbf{Z}} \stackrel{\text{def}}{=} \{\lambda^{Nk} \mid k \in \mathbf{Z}\}$, some $N \in \mathbf{N}$, $N \geq 2$. Let $B_1 = (A_1)_{\phi_1}(H)$. Note by Remark 4.3 that $\mathcal{M}$ is generated by B_1 and A_2 together with v . Let

$$\mathcal{N}_{[0,0]} = W^*(B_1 \cup A_2),$$

$$\mathcal{N}_{[0,n]} = W^*(\mathcal{N}_{[0,n-1]} \cup v^n \mathcal{N}_{[0,0]} (v^*)^n), \quad 1 \leq n \leq N-1.$$

Then

- (i) $h_n \mathcal{N}_{[0,n-1]} h_n$ and $v^n \mathcal{N}_{[0,0]} (v^*)^n$ are free in $(h_n \mathcal{M} h_n, \lambda^{-n} \phi|_{h_n \mathcal{M} h_n})$ with amalgamation over $h_n B_1 h_n$, $\forall 1 \leq n \leq N-1$, where $h_n = v^n (v^*)^n$;
- (ii) if also $\lambda^{\mathbf{Z}} \cap \Gamma' = \lambda^{N\mathbf{Z}}$, where Γ' is the subgroup of Γ generated by H together with Γ_2 , then $\mathcal{N}_{[0,N-1]} = \mathcal{M}(\Gamma')$.

Proof This is a modulo N version of the proof of Lemma 6.1. Because $v^N \in B_1$ and, e.g. for $1 \leq n \leq N-1$,

$$(v^*)^n B_1 = (v^*)^n (v^*)^{N-n} v^{N-n} B_1 (v^*)^{N-n} v^{N-n} = (v^*)^N (v^{N-n} B_1 (v^*)^{N-n}) v^{N-n} \subseteq B_1 v^{N-n}, \quad (16)$$

we have that $\text{span}(B_1 \cup \bigcup_{1 \leq n \leq N-1} B_1 v^n)$ is a s.o.-dense $*$ -subalgebra of A_1 , hence, letting

$$\Omega = \Lambda^\circ(B_1^\circ \cup \bigcup_{1 \leq n \leq N-1} B_1 v^n, A_2^\circ),$$

we have that $\text{span}(\Omega \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{M}$. Consider an alternating product $x = a_1 a_2 \cdots a_n \in \Omega$ with each $a_j \in B_1^\circ \cup \bigcup_{1 \leq n \leq N-1} B_1 v^n \subseteq A_1^\circ$ or $a_j \in A_2^\circ$ and with $a_j \in A_1^\circ \iff a_{j+1} \in A_2^\circ$. We will want to keep track of the number of occurrences of v modulo N , hence let $l(x) = n$ be the length of x and for each $1 \leq i, j \leq n$ let

$$s(a_i) = \begin{cases} 0 & \text{if } a_i \in B_1^\circ \cup A_2^\circ \\ k & \text{if } a_i \in B_1 v^k \end{cases}$$

$$t_j(x) = \sum_{i=1}^j s(a_i) \in \mathbf{Z}/N\mathbf{Z}, \quad (17)$$

taking the sum in (17) in the group of integers modulo N . If $0 \in I \subseteq \mathbf{Z}/N\mathbf{Z}$ let Θ_I be the set of words $x \in \Omega$ such that $t_{l(x)}(x) = 0$ and $t_j(x) \in I \forall 1 \leq j \leq l(x)$.

Claim 6.2a For each $1 \leq n \leq N-1$, $\text{span}(\Theta_{[0,n]} \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{N}_{[0,n]}$.

Proof One shows that $\Theta_{[0,n]}$ is a $*$ -algebra using induction just like in the proof of Claim 6.1a, and that $\mathcal{N}_{[0,n]} = \overline{\text{span}}(\Theta_{[0,n]} \cup \{1\})$ using double induction just like in the proof of (15).

The proof of (i) is similar to the proof of 6.1(ii). It will suffice to show that $\phi(y) = 0$ whenever y is an alternating product in $\Theta_{[0,n-1]} \setminus B_1^\circ$ and $v^n (\Theta_{[0,0]} \setminus B_1^\circ) (v^*)^n$. But every element of $\Theta_{[0,n-1]} \setminus B_1^\circ$ begins with an element of either A_2° , B_1° or $B_1 v^k$, ($1 \leq k \leq n-1$), and ends with an element of either A_2° , B_1° or $B_1 v^{N-k}$, ($1 \leq k \leq n-1$). Since, for $1 \leq k \leq n-1$, we have $B_1 v^{N-k} v^n B_1 \subseteq B_1 v^{n-k} \subseteq A_1^\circ$ and (see (16)) $B_1 (v^*)^n B_1 v^k \subseteq B_1 v^{N-n+k} \subseteq A_1^\circ$, by regrouping and multiplying some neighbors we see that y is equal to an alternating product in A_1° and A_2° , hence $\phi(y) = 0$ by freeness.

In light of Claim 6.2a, showing (ii) is the same as showing $\mathcal{M}_\phi(\Gamma') = \overline{\text{span}}(\Theta_{[0,N-1]} \cup \{1\})$. Again, note that $\Theta_{[0,N-1]}$ is the set of words, $x \in \Omega$ such that $t_{l(x)}(x) = 0$. Let E denote the projection from $\mathcal{M}$ onto $\mathcal{M}_\phi(\Gamma')$, obtained from Lemma 4.5. As in the proof of 6.1(iii), we see that every $x \in \Omega$ is the s.o.-limit of a bounded sequence in

$\text{span}(\bigcup_{\gamma \in \lambda^{k(x)} + \Gamma'} \mathcal{M}_\phi(\gamma))$, where $k(x) = -t_{l(x)}(x)$. Hence $t_{l(x)}(x) = 0$ implies $x \in \mathcal{M}_\phi(\Gamma')$ and $t_{l(x)}(x) \in \{1, \dots, N-1\}$ implies $x \perp \mathcal{M}_\phi(\Gamma')$, hence $E(x) = 0$. Thus $\mathcal{M}_\phi(\Gamma') \supseteq \overline{\text{span}}(\Theta_{[0, N-1]} \cup \{1\})$. To prove the reverse inclusion, let $y \in \mathcal{M}_\phi(\Gamma')$ and let $(y_n)_{n=1}^\infty$ be a bounded sequence in $\text{span}(\Omega \cup \{1\})$ whose s.o.-limit is y . Then $(E(y_n))_{n=1}^\infty$ is a bounded sequence in $\text{span}(\Theta_{[0, N-1]} \cup \{1\})$ whose s.o.-limit is $E(y) = y$. Hence $y \in \overline{\text{span}}(\Theta_{[0, N-1]} \cup \{1\})$, proving (ii). $\square$

Lemma 6.3 *Let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$, Γ_ι , $\Gamma_{\iota,0}$ and Γ be as in Lemma 6.1. Suppose $\lambda \in \Gamma_{1,0} \cap \Gamma_{2,0}$, $0 < \lambda < 1$, for each $\iota = 1, 2$ there is an isometry $v_\iota \in (A_\iota)_{\phi_\iota}(\lambda^{-1})$ and $\lambda^{\mathbf{Z}}$ is complemented in Γ_ι , say $\Gamma_\iota = H_\iota \oplus \lambda^{\mathbf{Z}}$. Let $B_\iota = (A_\iota)_{\phi_\iota}(H_\iota)$. Note by Remark 4.3 that $\mathcal{M}$ is generated by B_1 and B_2 together with v_1 and v_2 . For each $n \geq 1$, let $h_{\iota,n} = v_\iota^n (v_\iota^*)^n$. Then there is a partial isometry $y_n \in W^*(\{1, h_{1,n}, h_{2,n}\})$ such that $y_n^* y_n = h_{1,n}$ and $y_n y_n^* = h_{2,n}$. Let $u_n = y_n^* v_2^n (v_1^*)^n$. Let*

$$\begin{aligned} \mathcal{N}_{[0,0]} &= W^*(B_1 \cup B_2), \\ \mathcal{N}_{[0,1]} &= W^*(\mathcal{N}_{[0,0]} \cup \{u_1\}), \\ \mathcal{N}_{[-n,1]} &= W^*(v_1^* \mathcal{N}_{[-n+1,1]} v_1 \cup \mathcal{N}_{[0,1]}), \quad n \geq 1, \\ \mathcal{N}_{(-\infty,1]} &= W^*(\bigcup_{n \geq 0} \mathcal{N}_{[-n,1]}), \\ \mathcal{N}_{(-\infty,n]} &= W^*(\mathcal{N}_{(-\infty,n-1]} \cup \{u_n\}), \quad n \geq 2, \\ \mathcal{N}_{(-\infty,\infty)} &= W^*(\bigcup_{n \geq 1} \mathcal{N}_{(-\infty,n]}). \end{aligned}$$

Then

- (i) u_1 is a Haar unitary and $(h_{1,1} \mathcal{N}_{[0,0]} h_{1,1}, \{u_1\})$ is $*$ -free in $(h_{1,1} \mathcal{M} h_{1,1}, \lambda^{-1} \phi|_{h_{1,1} \mathcal{M} h_{1,1}})$;
- (ii) $v_1^* \mathcal{N}_{[-n+1,1]} v_1$ and $\mathcal{N}_{[0,1]}$ are free with amalgamation over $\mathcal{N}_{[0,0]}$ in $(\mathcal{M}, \phi)$, $\forall n \geq 1$;
- (iii) u_n is a Haar unitary and $(h_{1,n} \mathcal{N}_{(-\infty,n-1]} h_{1,n}, \{u_n\})$ is $*$ -free in $(h_{1,n} \mathcal{M} h_{1,n}, \lambda^{-n} \phi|_{h_{1,n} \mathcal{M} h_{1,n}})$, $\forall n \geq 2$;
- (iv) if $\lambda^{\mathbf{Z}} \cap \Gamma' = \{1\}$, where Γ' is the subgroup of Γ generated by H_1 together with H_2 , then $\mathcal{N}_{(-\infty,\infty)} = \mathcal{M}_\phi(\Gamma')$.

Proof Since $h_{1,n}$ and $h_{2,n}$ are free projections having the same weight under ϕ , the existence of the partial isometry y_n is easily seen from Theorem 1.1 of Dykema [1993(b)]. Since $\text{span}(B_\iota \cup \bigcup_{n \geq 1} (B_\iota v_\iota^n \cup (v_\iota^*)^n B_\iota))$ is a s.o.-dense $*$ -subalgebra of A_ι , letting

$$\Omega = \Lambda^\circ \left(B_1^\circ \cup \bigcup_{n \geq 1} (B_1 v_1^n \cup (v_1^*)^n B_1), B_2^\circ \cup \bigcup_{n \geq 1} (B_2 v_2^n \cup (v_2^*)^n B_2) \right),$$

we have that $\text{span}(\Omega \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{M}$. Consider a word $x = a_1 a_2 \cdots a_n$ belonging to Ω , such that $a_j \in B_{\iota_j}^\circ \cup \bigcup_{n \in \mathbf{N}} (B_{\iota_j} v_{\iota_j}^n \cup (v_{\iota_j}^*)^n B_{\iota_j}) \subseteq A_{\iota_j}^\circ$, with $\iota_j \neq \iota_{j+1}$. We will want to keep track of the occurrences of v_ι and v_ι^* , hence let $l(x) = n$ be the length of x and for each $1 \leq i, j \leq n$ let

$$\begin{aligned} s(a_i) &= \begin{cases} 0 & \text{if } a_i \in B_1^\circ \cup B_2^\circ \\ k & \text{if } a_i \in B_1 v_1^k \cup B_2 v_2^k \\ -k & \text{if } a_i \in (v_1^*)^k B_1 \cup (v_2^*)^k B_2, \end{cases} \\ t_j(x) &= \sum_{i=1}^j s(a_i). \end{aligned}$$

If $0 \in I \subseteq \mathbf{Z}$ let Θ_I be the set of words $x \in \Omega$ such that $t_{l(x)}(x) = 0$ and $t_j(x) \in I \forall 1 \leq j \leq l(x)$.

Claim 6.3a *If I is of the form $[0, 0]$, $[-n, 1]$ for $n \in \{0\} \cup \mathbf{N}$, $(-\infty, n]$ for $n \in \mathbf{N}$ or $(-\infty, \infty)$, then $\text{span}(\Theta_I \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{N}_I$.*

Proof One first proves that $\text{span}(\Theta_I \cup \{1\})$ is a $*$ -algebra like in Claim 6.1a. It is clear that $\mathcal{N}_{[0,0]} = \overline{\text{span}}(\Theta_{[0,0]} \cup \{1\})$. Let us prove that

$$\mathcal{N}_{[0,1]} = \overline{\text{span}}(\Theta_{[0,1]} \cup \{1\}). \quad (18)$$

Since $u_1 \in \overline{\text{span}}\Theta_{[0,1]}$, (which is seen by writing y_1 as the s.o.-limit of a bounded sequence in $\text{span}(\Theta_{[0,0]} \cup \{1\})$), the inclusion $\subseteq$ in (18) is clear. For the reverse inclusion, since $v_i^* v_i = 1$, it will suffice to show that $v_1 a v_1^*$, $v_1 a v_2^*$, $v_2 a v_2^* \in \mathcal{N}_{[0,1]}$ whenever $a \in B_1 \cup B_2$. If $a \in B_1$ then

$$\begin{aligned} v_1 a v_1^* &\in B_1 \subseteq \mathcal{N}_{[0,1]}, \\ v_1 a v_2^* &= (v_1 a v_1^*)(v_1 v_2^* y_1) y_1^* \in B_1 u_1^* y_1 \subseteq \mathcal{N}_{[0,1]}, \\ v_2 a v_2^* &= y_1 (y_1^* v_2 v_1^*) (v_1 a v_1^*) (v_1 v_2^* y_1) y_1^* \in y_1 u_1 B_1 u_1^* y_1^* \subseteq \mathcal{N}_{[0,1]}, \end{aligned} \quad (19)$$

and if $a \in B_2$ then

$$\begin{aligned} v_1 a v_1^* &= (v_1 v_2^* y) y^* (v_2 a v_2^*) y (y^* v_2 v_1^*) \in u_1^* y_1^* B_2 y_1 u_1 \subseteq \mathcal{N}_{[0,1]} \\ v_1 a v_2^* &= (v_1 v_2^* y) y^* (v_2 a v_2^*) \in u_1^* y_1^* B_2 \subseteq \mathcal{N}_{[0,1]} \\ v_2 a v_2^* &\in B_2 \subseteq \mathcal{N}_{[0,1]}. \end{aligned} \quad (20)$$

This proves (18).

Let us prove by induction on $n \geq 0$ that

$$\mathcal{N}_{[-n,1]} = \overline{\text{span}}(\Theta_{[-n,1]} \cup \{1\}). \quad (21)$$

The case $n = 0$ was just proved. Suppose $n \geq 1$. From the fact that $v_1^* \Theta_{[-n+1,1]} v_1 = \Theta_{[-n,0]}$ and from inductive hypothesis, one sees that $\subseteq$ holds in (21). To show the reverse inclusion, note that every element of $\Theta_{[-n,1]}$ equals a product of elements from $\Theta_{[-n,0]}$ and $\Theta_{[0,1]}$.

Now let us prove by induction on $n \geq 1$ that

$$\mathcal{N}_{(-\infty,n]} = \overline{\text{span}}(\Theta_{(-\infty,n]} \cup \{1\}). \quad (22)$$

The case $n = 1$ is true by taking the inductive limit of (21). For $n \geq 2$, the inclusion $\subseteq$ holds in (22) by inductive hypothesis and the fact (seen by writing y_n as a s.o.-limit of a bounded sequence in $\text{span}(\Theta_{[0,0]} \cup \{1\})$) that $u_n \subseteq \overline{\text{span}}\Theta_{[0,n]}$. For the reverse inclusion, analogous to the proof of (15), we show that every element of $\Theta_{(-\infty,n]}$ is a product of elements of $\Theta_{(-\infty,n-1]}$ and elements of the form $v_{\iota_1}^n \Theta_{[0,0]} (v_{\iota_2}^*)^n$, ($\iota_1, \iota_2 \in \{1, 2\}$). By inductive hypothesis, the former are in $\mathcal{N}_{(-\infty,n-1]}$, and the latter are shown to be in $\mathcal{N}_{(-\infty,n]}$ using analogues of (19) and (20) for u_n .

Finally, note that

$$\mathcal{N}_{(-\infty,\infty)} = \overline{\text{span}}(\Theta_{(-\infty,\infty)} \cup \{1\})$$

follows by taking the inductive limit of (22). Hence Claim 6.3a is proved.

Continuing with the proof of Lemma 6.3, we will now prove (i) and (iii) simultaneously. For consistency of notation, let $\mathcal{N}_{(-\infty,0]} = \overline{\text{span}}(\Theta_{(-\infty,0]} \cup \{1\})$, and note that $\mathcal{N}_{[0,0]} \subseteq \mathcal{N}_{(-\infty,0]}$. (We don't need to show that $\mathcal{N}_{(-\infty,0]}$ is an algebra; we only care that it contains $\mathcal{N}_{[0,0]}$.) We will thus show for every $n \geq 1$ that $\phi(z) = 0$ whenever z is

an alternating product in $(h_{1,n}\mathcal{N}_{(-\infty,n-1]}h_{1,n})^\circ$ and $\{u_n^k \mid k \in \mathbf{N}\} \cup \{(u_n^*)^k \mid k \in \mathbf{N}\}$. Rewriting $u_n = y_n^*v_2^n(v_1^*)^n$ and $u_n^* = v_1^n(v_2^*)^ny_n$ and regrouping, we see that z equals an alternating product, call it z' , in

$$(h_{1,n}\mathcal{N}_{(-\infty,n-1]}h_{1,n})^\circ \cup (h_{1,n}\mathcal{N}_{(-\infty,n-1]}h_{2,n}) \cup (h_{2,n}\mathcal{N}_{(-\infty,n-1]}h_{1,n}) \cup (h_{2,n}\mathcal{N}_{(-\infty,n-1]}h_{2,n})^\circ$$

and $\{v_2^n(v_1^*)^n, v_1^n(v_2^*)^n\}$, where every letter that has

$$v_2^n(v_1^*)^n \text{ on the left and } v_1^n(v_2^*)^n \text{ on the right belongs to } (h_{1,n}\mathcal{N}_{(-\infty,n-1]}h_{1,n})^\circ$$

$$v_1^n(v_2^*)^n \text{ on the left and } v_2^n(v_1^*)^n \text{ on the right belongs to } (h_{2,n}\mathcal{N}_{(-\infty,n-1]}h_{2,n})^\circ.$$

From Claim 6.3a and Kaplansky's density theorem, we know that every element of $\mathcal{N}_{(-\infty,n-1]}$ is the s.o.-limit of a bounded sequence in $\text{span}(\Theta_{(-\infty,n-1]} \cup \{1\})$. Using now the conditional expectation from $\mathcal{M}$ onto B_ι , gotten from combining Corollary 3.4 and Lemma 4.5, we see that every element of $(h_{\iota,n}\mathcal{N}_{(-\infty,n-1]}h_{\iota,n})^\circ$ is the s.o.-limit of a bounded sequence in $\text{span}((\Theta_{(-\infty,n-1]} \setminus B_\iota^\circ) \cup (h_{\iota,n}B_\iota h_{\iota,n})^\circ)$. Hence, to show that $\phi(z') = 0$ it will suffice to show that $\phi(z'') = 0$ whenever z'' is obtained from z' by replacing every letter of z' that

lies between $v_2^n(v_1^*)^n$ on the left and $v_1^n(v_2^*)^n$ on the right

$$\text{with an arbitrary element of } (\Theta_{(-\infty,n-1]} \setminus B_1^\circ) \cup (h_{1,n}B_1h_{1,n})^\circ$$

lies between $v_1^n(v_2^*)^n$ on the left and $v_2^n(v_1^*)^n$ on the right

$$\text{with an arbitrary element of } (\Theta_{(-\infty,n-1]} \setminus B_2^\circ) \cup (h_{2,n}B_2h_{2,n})^\circ,$$

and replacing every letter of z' not in one of the above two cases with an arbitrary element of $\Theta_{(-\infty,n-1]} \cup \{1\}$. Now regrouping and multiplying some neighbors in z'' and using that $(v_\iota^*)^n(h_{\iota,n}B_\iota h_{\iota,n})^\circ v_\iota^n \subseteq B_\iota^\circ$ and $v_\iota^n B_\iota \subseteq A_\iota^\circ$, we see that z'' is equal to an alternating product in A_1° and A_2° , hence by freeness $\phi(z'') = 0$. Thus (i) and (iii) are proved.

To prove part (ii), we must show that $\phi(z) = 0$ whenever z is an alternating product in $v_1^*\mathcal{N}_{[-n+1,1]}v_1 \ominus \mathcal{N}_{[0,0]}$ and $\mathcal{N}_{[0,1]} \ominus \mathcal{N}_{[0,0]}$. Let E be the conditional expectation from $\mathcal{M}$ onto $\mathcal{N}_{[0,0]}$ gotten from Lemma 3.3 and Lemma 4.5. If $x \in v_1^*\mathcal{N}_{[-n+1,1]}v_1$, then by Claim 6.3a and Kaplansky's density theorem, x is the s.o.-limit of a bounded sequence, $(x_n)_{n=1}^\infty$, in $\text{span}(v_1^*\Theta_{[-n+1,1]}v_1 \cup \{1\}) = \text{span}(\Theta_{[-n,0]} \cup \{1\})$. If also $x \perp \mathcal{N}_{[0,0]}$, then $E(x) = 0$, so $x_n - E(x_n)$ converges strongly to x as $n \rightarrow \infty$, and $x_n - E(x_n) \in \text{span}(\Theta_{[-n,0]} \setminus \Theta_{[0,0]})$. Hence every element of $v_1^*\mathcal{N}_{[-n+1,1]}v_1 \ominus \mathcal{N}_{[0,0]}$ is the s.o.-limit of a bounded sequence in $\text{span}(\Theta_{[-n,0]} \setminus \Theta_{[0,0]})$. Similarly, every element of $\mathcal{N}_{[0,1]} \ominus \mathcal{N}_{[0,0]}$ is the s.o.-limit of a bounded sequence in $\text{span}(\Theta_{[0,1]} \setminus \Theta_{[0,0]})$. Hence it will suffice to show that $\phi(z') = 0$ whenever z' is an alternating product in $\Theta_{[-n,0]} \setminus \Theta_{[0,0]}$ and $\Theta_{[0,1]} \setminus \Theta_{[0,0]}$. However, note that $\Theta_{[0,0]}(\Theta_{[-n,0]} \setminus \Theta_{[0,0]})\Theta_{[0,0]} \subseteq \text{span}(\Theta_{[-n,0]} \setminus \Theta_{[0,0]})$. Hence from every letter of z' that belongs to $\Theta_{[-n,0]} \setminus \Theta_{[0,0]}$, we can peel off any leading or trailing letters from B_1° or B_2° and glue them onto the adjoining letters of z' , (except if we're at the beginning or the end of z'). Therefore, letting Ψ be the set of words $x = a_1 \cdots a_n \in \Theta_{[0,1]}$ such that $a_1 \in B_\iota v_\iota$, some $\iota = 1, 2$ and $a_n \in v_\iota^* B_\iota$, some $\iota = 1, 2$, it will suffice to show that $\phi(fz''g) = 0$ whenever z'' is an alternating product in $\Theta_{[-n,0]} \setminus \Theta_{[0,0]}$ and Ψ , and whenever $f = 1$ (respectively $g = 1$) if the first letter (respectively last letter) of z'' is from $\Theta_{[-n,0]} \setminus \Theta_{[0,0]}$ and f (respectively g) is in $\Theta_{[0,0]} \cup \{1\}$ if the first letter (respectively last letter) of z'' is in Ψ . By regrouping and multiplying some neighbors, $fz''g$ is seen to be an alternating product in A_1° and A_2° , hence $\phi(fz''g) = 0$ by freeness and (ii) is proved.

In light of the Claim 6.3a, part (iv) is proved just like 6.1(iii) was proved. $\square$

Lemma 6.4 *Let $(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$, Γ_ι , $\Gamma_{\iota,0}$ and Γ be as in Lemma 6.1. Suppose $\lambda \in \Gamma_{1,0} \cap \Gamma_{2,0}$, $0 < \lambda < 1$, for each $\iota = 1, 2$ there is an isometry $v_\iota \in (A_\iota)_{\phi_\iota}(\lambda^{-1})$ and there is a subgroup $H_\iota \subseteq \Gamma_\iota$ that together with λ generates Γ_ι . Suppose $\lambda^{\mathbf{Z}} \cap H_1 = \lambda^{N\mathbf{Z}} = \lambda^{\mathbf{Z}} \cap H_2$, some $N \in \mathbf{N}$, $N \geq 2$. Let $B_\iota = (A_\iota)_{\phi_\iota}(H_\iota)$. Note by Remark 4.3 that $\mathcal{M}$ is generated by B_1 and B_2 together with v_1 and v_2 . For each $n \geq 1$, let $h_{\iota,n} = v_\iota^n (v_\iota^*)^n$. Then there is a partial isometry $y_n \in W^*(\{1, h_{1,n}, h_{2,n}\})$ such that $y_n^* y_n = h_{1,n}$ and $y_n y_n^* = h_{2,n}$. Let $u_n = y_n^* v_2^n (v_1^*)^n$. Let*

$$\mathcal{N}_{[0,0]} = W^*(B_1 \cup B_2),$$

$$\mathcal{N}_{[0,n]} = W^*(\mathcal{N}_{[0,n-1]} \cup \{u_n\}), \quad 1 \leq n \leq N-1.$$

Then

- (i) $\{u_n\}$ is a Haar unitary and $(h_{1,n} \mathcal{N}_{[0,n-1]} h_{1,n}, \{u_n\})$ is $*$ -free in $(h_{1,n} \mathcal{M} h_{1,n}, \lambda^{-n} \phi|_{h_{1,n} \mathcal{M} h_{1,n}})$, $\forall 1 \leq n \leq N-1$;
- (ii) if $\lambda^{\mathbf{Z}} \cap \Gamma' = \lambda^{N\mathbf{Z}}$, where Γ' is the subgroup of Γ generated by H_1 and H_2 , then $\mathcal{M}_\phi(\Gamma') = \mathcal{N}_{[0,N-1]}$.

Proof This is a modulo N version of the proof of Lemma 6.3. As in the proof of Lemma 6.2, we have that $\text{span}(B_\iota \cup \bigcup_{1 \leq n \leq N-1} B_\iota v_\iota^n)$ is a s.o.-dense $*$ -subalgebra of A_ι , hence, letting

$$\Omega = \Lambda^\circ(B_1^\circ \cup \bigcup_{1 \leq n \leq N-1} B_1 v_1^n, B_2^\circ \cup \bigcup_{1 \leq n \leq N-1} B_2 v_2^n),$$

we have that $\text{span}(\Omega \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{M}$. Consider a word $x = a_1 a_2 \cdots a_n \in \Omega$ with each $a_j \in B_{\iota_j}^\circ \cup \bigcup_{1 \leq n \leq N-1} B_{\iota_j} v_{\iota_j}^n \subseteq A_{\iota_j}^\circ$ and with $\iota_j \neq \iota_{j+1}$. We will want to keep track of the number of occurrences of v_1 and v_2 modulo N , hence let $l(x) = n$ be the length of x and for each $1 \leq i, j \leq n$ let

$$\begin{aligned} s(a_i) &= \begin{cases} 0 & \text{if } a_i \in B_1^\circ \cup B_2^\circ \\ k & \text{if } a_i \in B_1 v_1^k \cup B_2 v_2^k \end{cases} \\ t_j(x) &= \sum_{i=1}^j s(a_i) \in \mathbf{Z}/N\mathbf{Z}, \end{aligned} \tag{23}$$

taking the sum in (23) in the group of integers modulo N . If $0 \in I \subseteq \mathbf{Z}/N\mathbf{Z}$ let Θ_I be the set of words $x \in \Omega$ such that $t_{l(x)}(x) = 0$ and $t_j(x) \in I \forall 1 \leq j \leq l(x)$.

Claim 6.4a *For each $1 \leq n \leq N-1$, $\text{span}(\Theta_{[0,n]} \cup \{1\})$ is a s.o.-dense $*$ -subalgebra of $\mathcal{N}_{[0,n]}$.*

Proof One shows that $\Theta_{[0,n]}$ is a $*$ -algebra using induction just like in the proof of Claim 6.1a, and that $\mathcal{N}_{[0,n]} = \overline{\text{span}}(\Theta_{[0,n]} \cup \{1\})$ just like in the proofs of (18) and (22).

Proving (i) is like proving 6.3(i) and (iii).

In light of the Claim 6.4a, part (ii) is proved just like 6.1(iii) and 6.2(ii) were proved. We omit the details. $\square$

Proposition 6.5 *Let A_1 be a separable, type III factor with extremal almost periodic state ϕ_1 and suppose the centralizer of ϕ_1 is either the hyperfinite II_1 -factor, R , or $L(F_\infty)$. Let A_2 be a separable von Neumann algebra with normal, faithful state ϕ_2 and suppose either*

(diff) ϕ_2 is a trace, A_2 is either $L(\mathbf{Z})$ or an interpolated free group factor,

(fin) ϕ_2 is a trace and $A_2 \neq \mathbf{C}$ is the direct sum of a finite dimensional algebra, a hyperfinite algebra and (possibly infinitely many) interpolated free group factors

or

(extr) A_2 is a type III factor and ϕ_2 is an extremal almost periodic state whose centralizer is either R or $L(F_\infty)$.

Let

$$(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2).$$

Then $\mathcal{M}_\phi \cong L(F_\infty)$. Furthermore, the inclusion $Q \hookrightarrow \mathcal{M}_\phi$ is a standard embedding, where $Q = W^*((A_1)_{\phi_1} \cup (A_2)_{\phi_2}) \cong L(F_t)$. Thus $\mathcal{M}$ is a type III factor and ϕ is an extremal almost periodic state whose point spectrum $Sd(\phi)$ is the subgroup of $\mathbf{R}_+^*$ generated by $Sd(\phi_1) \cup Sd(\phi_2)$.

Note that part of this proposition is Theorem 2.

Proof of Proposition 6.5 We need only show that $\mathcal{M}_\phi \cong L(F_\infty)$ and $Q \hookrightarrow \mathcal{M}_\phi$ is a standard embedding. These assertions will be proved by induction in several steps, corresponding to several cases. First, we show that

$$(\text{diff}) \implies (\text{fin}),$$

or more precisely that whenever the proposition has been proved in case (diff) for a given value of $Sd(\phi_1)$, then the conclusions of the proposition hold also in the case (fin) for that same value of $Sd(\phi_1)$. Indeed, in case (fin), by Dykema [1993(b)] there is n large enough so that if $h_1, \dots, h_n \in (A_1)_{\phi_1}$ are orthogonal projections, $\phi_1(h_j) = 1/n \forall j$, then

$$\mathcal{P}_0 \stackrel{\text{def}}{=} W^*({h_1, \dots, h_n} \cup A_2)$$

is an interpolated free group factor. For $2 \leq j \leq n$ let $v_j \in (A_1)_{\phi_1}$ be a partial isometry such that $v_j^* v_j = h_1$ and $v_j v_j^* = h_j$. Then $A_1 = W^*(h_1 A_1 h_1 \cup \{v_2, \dots, v_n\})$. Let

$$\mathcal{P}_1 = W^*(\mathcal{P}_0 \cup h_1 A_1 h_1)$$

$$\mathcal{P}_j = W^*(\mathcal{P}_{j-1} \cup \{v_j\}) \quad 2 \leq j \leq n.$$

Thus $\mathcal{P}_n = \mathcal{M}$. We have by Proposition 5.1 that $h_1 \mathcal{P}_1 h_1$ is freely generated by $h_1 \mathcal{P}_0 h_1$ and $h_1 A_1 h_1$, so by the proposition in case (diff), and since h_1 has central support equal to 1, $\mathcal{P}_1$ is a type III factor, $\phi|_{\mathcal{P}_1}$ is an extremal almost periodic state with centralizer isomorphic to $L(F_\infty)$ and $Sd(\phi|_{\mathcal{P}_1}) = Sd(\phi_1)$. In exactly the same way, using Lemma 5.4, we have by induction on $2 \leq j \leq n$ that $\mathcal{P}_j$ is a type III factor, $\phi|_{\mathcal{P}_j}$ is an extremal almost periodic state with centralizer isomorphic to $L(F_\infty)$ and $Sd(\phi|_{\mathcal{P}_j}) = Sd(\phi_1)$. It only remains to show that $\mathcal{Q} \hookrightarrow \mathcal{M}_\phi$ is a standard embedding. We have that $W^*(h_1 \mathcal{P}_0 h_1 \cup h_1 (A_1)_{\phi_1} h_1) \hookrightarrow (h_1 \mathcal{P}_1 h_1)_\phi$ is a standard embedding, and by choosing x to lie in $\mathcal{P}_0$ for the applications of Lemma 5.4 to each inclusion $h_1 \mathcal{P}_{j-1} h_1 \hookrightarrow h_1 \mathcal{P}_j h_1$, we see that $h_1 (W^*(h_1 A_1 h_1 \cup \{v_2, \dots, v_j\}) \cup A_2) h_1 \hookrightarrow h_1 (\mathcal{P}_j)_\phi h_1$ is a standard embedding $\forall 2 \leq j \leq n$, hence at $j = n$ we have that $\mathcal{Q} \hookrightarrow \mathcal{M}_\phi$ is a standard embedding. Thus the conclusions of the proposition hold also in case (fin).

Henceforth, whenever $D \subseteq \mathcal{M}$ is a von Neumann subalgebra, by D_ϕ we will denote the centralizer of $\phi|_D$.

We begin by showing that the conclusions of the proposition hold in the following cases for $1 \leq m \leq \infty$.

Case I(diff)_m ϕ_2 is a trace, A_2 is $L(\mathbf{Z})$ or an interpolated free group factor and $Sd(\phi_1)$ is a free abelian group of rank m .

Case I(extr)_m A_2 is a type III factor, ϕ_2 is an extremal almost periodic state, $Sd(\phi_2) = Sd(\phi_1)$ and $Sd(\phi_1)$ is a free abelian group of rank m .

Our strategy will be to prove

- (a) I(diff)₁ and
I(diff)_{m-1} $\implies$ I(diff)_m, ($2 \leq m < \infty$)
- (b) I(extr)₁ and
I(diff)_{m-1} + I(extr)_{m-1} $\implies$ I(extr)_m, ($2 \leq m < \infty$)
- (c) I(diff)_∞ and I(extr)_∞.

Let us begin by proving (a), *i.e.* I(diff)_m, $1 \leq m < \infty$. We will use Lemma 6.1 with $Sd(\phi_1) = \lambda^{\mathbf{Z}} \oplus H$, for H a free abelian group of rank $m - 1$, ($H = \{1\}$ when $m = 1$). Then also part 6.1(iii) applies, and hence $\mathcal{M}_\phi = (\mathcal{N}_{(-\infty, \infty)})_\phi$. We will show by induction on $n \geq 1$ that

- (a1) $\mathcal{N}_{[-n, 0]}$ is a factor, $\phi|_{\mathcal{N}_{[-n, 0]}}$ is an extremal almost periodic state with $Sd(\phi|_{\mathcal{N}_{[-n, 0]}}) = H$.
- (a2) B_1 is freely subcomplemented in $\mathcal{N}_{[-n, 0]}$ by a finite family $(D_\iota)_{\iota \in I}$ of algebras, where each D_ι is either $L(\mathbf{Z})$ or $L(F_t)$, $1 < t \leq \infty$ with $\phi|_{D_\iota}$ being a trace.
- (a3) $\mathcal{N}_{[-n+1, 0]}$ is freely subcomplemented in $\mathcal{N}_{[-n, 0]}$ by a finite family of algebras, each of which is of the form described in (a2).
- (a4) $(\mathcal{N}_{[-n, 0]})_\phi \cong L(F_{t_n})$ with $t_n = \infty$ (or if $m = 1$ possibly $t_n < \infty$ but $t_n \nearrow \infty$).
- (a5) the inclusion $(\mathcal{N}_{[-n+1, 0]})_\phi \hookrightarrow (\mathcal{N}_{[-n, 0]})_\phi$ is a standard embedding.
- (a6) the inclusion $\mathcal{Q} \hookrightarrow (\mathcal{N}_{[-n, 0]})_\phi$ is a standard embedding.

By definition, B_1 is freely complemented in $\mathcal{N}_{[0, 0]}$ by A_2 , hence by I(diff)_{m-1} (or by Dykema [1993(b)] in the case $m = 1$), we have that (a1), (a2), (a4) and (a6) hold for $n = 0$. Now suppose $n \geq 1$. We use 6.1(i) and we want to apply Lemma 2.3. We need that B_1 is freely subcomplemented in $v^* \mathcal{N}_{[-n+1, 0]} v$, and since $v^* B_1 v = B_1$, this is equivalent to $h_1 B_1 h_1$ being freely subcomplemented in $h_1 \mathcal{N}_{[-n+1, 0]} h_1$. From (a2)_{n-1} and from Lemma 5.6, making use of I(fin)_{m-1}, we see that $h_1 B_1 h_1$ is freely subcomplemented in $h_1 \mathcal{N}_{[-n+1, 0]} h_1$ by algebras of the form described in (a2). Hence, by 2.3, we have that (a2)_n holds. If $n = 1$ then also by 2.3, (a3)_n holds. If $n \geq 2$ then by (a2)_{n-2} and (a3)_{n-1}, applying 5.6 twice, we have that B_1 is freely subcomplemented in $v^* \mathcal{N}_{[-n+2, 0]} v$, which is in turn freely subcomplemented in $v^* \mathcal{N}_{[-n+1, 0]} v$ by $(D_\iota)_{\iota \in I}$, all by algebras of the form described in (a2). Thus, using the idea of the proof of 2.3, we see that $\mathcal{N}_{[-n+1, 0]} = W^*(v^* \mathcal{N}_{[-n+2, 0]} v \cup \mathcal{N}_{[0, 0]})$ is freely subcomplemented in $\mathcal{N}_{[-n, 0]}$ by $(D_\iota)_{\iota \in I}$, *i.e.* (a3)_n holds. Now (a4)_n, (a5)_n and (a6)_n all follow from (a3)_n using I(diff)_{m-1} and I(extr)_{m-1}.

From these we conclude that $\mathcal{N}_{(-\infty, 0]}$ is a factor, $\phi|_{\mathcal{N}_{(-\infty, 0]}}$ is an extremal almost periodic state with $Sd(\phi|_{\mathcal{N}_{(-\infty, 0]}}) = H$ and $(\mathcal{N}_{(-\infty, 0]})_\phi \cong L(F_\infty)$. Moreover, we have that $\mathcal{Q} \hookrightarrow (\mathcal{N}_{(-\infty, 0]})_\phi$ is a standard embedding and that B_1 is freely subcomplemented in $\mathcal{N}_{(-\infty, 0]}$ by a family of algebras of the form described in (a2). We can then use 6.1(ii) to prove by induction on $n \geq 1$ that

- (a7) $\mathcal{N}_{(-\infty, n]}$ is a factor, $\phi|_{\mathcal{N}_{(-\infty, n]}}$ is an extremal almost periodic state with $Sd(\phi|_{\mathcal{N}_{(-\infty, n]}}) = H$ and $(\mathcal{N}_{(-\infty, n]})_\phi \cong L(F_\infty)$.
- (a8) $h_n B_1 h_n$ is freely subcomplemented in $h_n \mathcal{N}_{(-\infty, n]} h_n$ by a countable family of algebras, each of which is of the form described in (a2).
- (a9) $h_n \mathcal{N}_{(-\infty, n-1]} h_n$ is freely subcomplemented in $h_n \mathcal{N}_{(-\infty, n]} h_n$ by a finite family of algebras, each of which is of the form described in (a2).
- (a10) the inclusion $(\mathcal{N}_{(-\infty, n-1]})_\phi \hookrightarrow (\mathcal{N}_{(-\infty, n]})_\phi$ is a standard embedding.
- (a11) the inclusion $\mathcal{Q} \hookrightarrow (\mathcal{N}_{(-\infty, n]})_\phi$ is a standard embedding.

Hence by taking inductive limits we have that $\mathcal{N}_{(-\infty, \infty)}$ is a factor, $\phi|_{\mathcal{N}_{(-\infty, \infty)}}$ is extremal almost periodic with $Sd(\phi|_{\mathcal{N}_{(-\infty, \infty)}}) = H$ and $(\mathcal{N}_{(-\infty, \infty)})_\phi \cong L(F_\infty)$ and $\mathcal{Q} \hookrightarrow (\mathcal{N}_{(-\infty, \infty)})_\phi$ is a standard embedding. Thus part (a) is proved.

Now we prove (b), *i.e.* $I(\text{extr})_m$, $1 \leq m < \infty$. We will use Lemma 6.3 with $Sd(\phi_1) = Sd(\phi_2) = \lambda^{\mathbf{Z}} \oplus H$, and note that 6.3(iv) applies, so we need only show that $(\mathcal{N}_{(-\infty, \infty)})_\phi \cong L(F_\infty)$ and $\mathcal{Q} \hookrightarrow (\mathcal{N}_{(-\infty, \infty)})_\phi$ is a standard embedding. We have by $I(\text{extr})_{m-1}$ (or by Dykema [1994(a)] if $m = 1$) that $\mathcal{N}_{[0, 0]}$ is a factor, $\phi|_{\mathcal{N}_{[0, 0]}}$ is an extremal almost periodic state, $Sd(\phi|_{\mathcal{N}_{[0, 0]}}) = H$, $(\mathcal{N}_{[0, 0]})_\phi \cong L(F_\infty)$ (or possibly $\cong L(F_2)$ if $m = 1$) and $\mathcal{Q} \hookrightarrow (\mathcal{N}_{[0, 0]})_\phi$ is a standard embedding (or $\mathcal{Q} = (\mathcal{N}_{[0, 0]})_\phi$ if $m = 1$). Now we will show by induction on $n \geq 1$ that

- (b1) $\mathcal{N}_{[-n, 1]}$ is a factor, $\phi|_{\mathcal{N}_{[-n, 1]}}$ is an extremal almost periodic state with $Sd(\phi|_{\mathcal{N}_{[-n, 1]}}) = H$.
- (b2) $\mathcal{N}_{[0, 1]}$ is freely subcomplemented in $\mathcal{N}_{[-n, 1]}$ by a finite family $(D_\iota)_{\iota \in I}$ of algebras, where each D_ι is either $L(\mathbf{Z})$, $L(F_t)$, $1 < t \leq \infty$ with $\phi|_{D_\iota}$ being a trace, or is a type III factor with $\phi|_{D_\iota}$ being an extremal almost periodic state having centralizer isomorphic to $L(F_\infty)$ and $Sd(\phi|_{D_\iota}) = H$.
- (b3) $\mathcal{N}_{[-n+1, 1]}$ is freely subcomplemented in $\mathcal{N}_{[-n, 1]}$ by a finite family of algebras, each of which is of the form described in (b2).
- (b4) $(\mathcal{N}_{[-n, 1]})_\phi \cong L(F_{t_n})$ with $t_n = \infty$ (or if $m = 1$ possibly $t_n < \infty$ but $t_n \nearrow \infty$).
- (b5) the inclusion $(\mathcal{N}_{[-n+1, 1]})_\phi \hookrightarrow (\mathcal{N}_{[-n, 1]})_\phi$ is a standard embedding.
- (b6) the inclusion $\mathcal{Q} \hookrightarrow (\mathcal{N}_{[-n, 1]})_\phi$ is a standard embedding.

By 6.3(i) and the proposition in case $I(\text{diff})_m$, we see that (b1), (b4) and (b6) hold for $n = 0$. Now suppose $n \geq 1$. We want to use 6.3(ii) and Proposition 2.3, and hence we must show that $\mathcal{N}_{[0, 0]}$ is freely subcomplemented in $v_1^* \mathcal{N}_{[-n+1, 1]} v_1$. This is equivalent to showing that $v_1 \mathcal{N}_{[0, 0]} v_1^*$ is freely subcomplemented in $h_1 \mathcal{N}_{[-n+1, 1]} h_1$. Applying Lemma 5.6 and using (b2) $_{n-1}$ if $n \geq 2$, it thus suffices that $v_1 \mathcal{N}_{[0, 0]} v_1^*$ be freely subcomplemented in $h_1 \mathcal{N}_{[0, 1]} h_1$. Now $v_1 \mathcal{N}_{[0, 0]} v_1^*$ is generated by $v_1 B_1 v_1^* = h_1 B_1 h_1$ and $v_1 B_2 v_1^* = u_1^* y_1^* B_2 y_1 u_1$ (with u_1 and y_1 as in 6.3). Consider

$$D = W^*(\{1\} \cup h_1 B_1 h_1 \cup h_2 B_2 h_2) = (\mathbf{C}(1 - h_1) \oplus h_1 B_1 h_1) * (\mathbf{C}(1 - h_2) \oplus h_2 B_2 h_2)$$

$$D \supseteq D_1 = (\mathbf{C}(1 - h_1) \oplus \mathbf{C} h_1) * (\mathbf{C}(1 - h_2) \oplus \mathbf{C} h_2).$$

Then by Lemma 5.3, $h_1 B_1 h_1$ and $y_1^* B_2 y_1$ are free and $W^*(h_1 B_1 h_1 \cup y_1^* B_2 y_1)$ is freely complemented in $h_1 D h_1$ by $h_1 D_1 h_1$, which by Theorem 1.1 of Dykema [1993(b)] is of the form $L(\mathbf{Z})$ or $L(\mathbf{Z}) \oplus \mathbf{C}$. Hence, adding partial

isometries from $(B_1)_{\phi_1}$ and $(B_2)_{\phi_2}$ like in the proof of Lemma 5.6, we see that $W^*(h_1 B_1 h_1 \cup y_1^* B_2 y_1)$ is freely subcomplemented in $h_1 \mathcal{N}_{[0,0]} h_1$ by a finite family $(D_\iota)_{\iota \in I}$ of algebras of the form $L(\mathbf{Z})$ or $L(\mathbf{Z}) \oplus \mathbf{C}$, and all associated projections can be chosen to lie in $h_1 (B_1)_\phi h_1$. Because $h_1 \mathcal{N}_{[0,0]} h_1$ and $\{u_1\}$ together freely generate $h_1 \mathcal{N}_{[0,1]} h_1$, by a free etymology argument it follows that $h_1 B_1 h_1, u_1^* y_1^* B_2 y_1 u_1, (D_\iota)_{\iota \in I}$ and $\{u_1\}$ together generate $h_1 \mathcal{N}_{[0,1]} h_1$, that $(h_1 B_1 h_1, u_1^* y_1^* B_2 y_1 u_1, \{u_1\})$ is $*$ -free in $h_1 \mathcal{N}_{[0,1]} h_1$ and that $W^*(h_1 B_1 h_1 \cup u_1^* y_1^* B_2 y_1 u_1 \cup \{u_1\})$ is freely subcomplemented in $h_1 \mathcal{N}_{[0,1]} h_1$ by $(D_\iota)_{\iota \in I}$; therefore $v_1 \mathcal{N}_{[0,0]} v_1^* = W^*(h_1 B_1 h_1 \cup u_1^* y_1^* B_2 y_1 u_1)$ is freely subcomplemented in $h_1 \mathcal{N}_{[0,1]} h_1$ by algebras of the form described in (a2). Now Proposition 2.3 can be used to show (b2) $_n$ and, in conjunction with (b3) $_{n-1}$ if $n \geq 2$, to show (b3) $_n$. Then (b4) $_n$, (b5) $_n$ and (b6) $_n$ follow as a matter of course.

Thus we see that $\mathcal{N}_{(-\infty, 1]}$ is a factor, $\phi|_{\mathcal{N}_{(-\infty, 1]}}$ is extremal almost periodic and has centralizer isomorphic to $L(F_\infty)$. Now by 6.3(iii) it is immediate that $\mathcal{N}_{(-\infty, 1]}$ is freely subcomplemented in $\mathcal{N}_{(-\infty, \infty)}$ by a countable family of algebras isomorphic to $L(\mathbf{Z})$ and from this (b) is easily proved.

Then (c) follows by taking inductive limits, and hence the proposition is proved in Case I(diff) and Case I(extr).

Next, consider

Case II ϕ_2 is a trace and A_2 is either $L(\mathbf{Z})$ or an interpolated free group factor.

Of course, if $Sd(\phi_1)$ is a free abelian group, then this has been proved as Case I(diff). Since every countable subgroup of $\mathbf{R}_+^*$ is the union of a chain of subgroups $H_1 \subseteq H_2 \subseteq \dots$ such that each H_j is a free abelian group and such that H_{j+1}/H_j is a finite cyclic group $\forall j$, one can prove the proposition in Case II using Case I(diff) and Lemma 6.2.

Thus all cases (diff) and (fin) have been proved. Consider now

Case III A_1 is a type III factor, ϕ_2 is an extremal almost periodic state and $Sd(\phi_1) = Sd(\phi_2)$.

This is easily proved using Case I(extr), the chain $H_1 \subseteq H_2 \subseteq \dots$ and Lemma 6.4.

It remains to show the case when A_1 and A_2 are type III factors but $Sd(\phi_1) \neq Sd(\phi_2)$. Let $H = Sd(\phi_1) \cap Sd(\phi_2)$ and let $H = K_1^{(\iota)} \subseteq K_2^{(\iota)} \subseteq \dots \subseteq Sd(\phi_\iota)$ ($\iota = 1, 2$) be a finite or infinite sequence of subgroups such that $Sd(\phi_\iota) = \bigcup_j K_j^{(\iota)}$ and $K_{j+1}^{(\iota)}/K_j^{(\iota)}$ is cyclic $\forall j$. If x is an element of $K_{j+1}^{(\iota)}$ that together with $K_j^{(\iota)}$ generates $K_{j+1}^{(\iota)}$, then $\mathbf{Z}x \cap K_j^{(\iota)} = \mathbf{Z}x \cap (K_j^{(\iota)} + Sd(\phi_{\iota'}))$ where $\iota' \neq \iota$. Hence by using Case III to get started, considering in turn $W^*((A_1)_{\phi_1}(K_j^{(1)}) \cup A_2(H))$ and $W^*(A_1 \cup (A_2)_{\phi_2}(K_j^{(2)}))$ and using Lemmas 6.1 and 6.2, one proves the proposition.

□

7. Free products of finite dimensional algebras.

Definition 7.1 For a finite dimensional von Neumann algebra A and a faithful positive linear functional ϕ on A , we define the nonnegative integer $\text{ntr}(A, \phi)$, which very roughly measures how far ϕ is from being a trace, as follows. If $A = M_n(\mathbf{C})$ and

$$\phi = \text{Tr} \left(\cdot \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix} \right) \quad (24)$$

then $\text{ntr}(M_n(\mathbf{C}), \phi) = 0$ if ϕ is a trace and $\text{ntr}(M_n(\mathbf{C}), \phi) = n$ otherwise. For a general finite dimensional A , we may write $A = \bigoplus_{j=1}^N A_j$ where each A_j is a matrix algebra and we define $\text{ntr}(A, \phi) = \sum_{j=1}^N \text{ntr}(A_j, \phi|_{A_j})$.

Remark 7.2 Let us take this opportunity to observe that the point spectrum of the modular operator, Δ_ϕ , of a faithful state ϕ on a finite dimensional algebra A is easily computed. Indeed, if $A = M_n(\mathbf{C})$ and ϕ is as in (24), then

$$\text{point spectrum}(\Delta_\phi) = \left\{ \frac{\lambda_i}{\lambda_j} \mid 1 \leq i, j \leq n \right\}$$

and for a faithful state on a general finite dimensional A , the point spectrum of Δ_ϕ is just the union of the sets of such ratios for all the simple summands of A .

Proposition 7.3 *Let*

$$(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$$

where A_1 is a type III factor, ϕ_1 is an extremal almost periodic state on A_1 with centralizer isomorphic to R or $L(F_\infty)$, A_2 is finite dimensional, $A_2 \neq \mathbf{C}$ and ϕ_2 is faithful on A_2 . Then $\mathcal{M}$ is a type III factor, ϕ is an extremal almost periodic state with centralizer isomorphic to $L(F_\infty)$ and $Sd(\phi)$ is the subgroup of $\mathbf{R}_+^*$ generated by

$$Sd(\phi_1) \cup (\text{point spectrum}(\Delta_{\phi_2})).$$

Furthermore, if $(A_1)_{\phi_1} \cong L(F_\infty)$ then the inclusion $(A_1)_{\phi_1} \hookrightarrow \mathcal{M}_\phi$ is a standard embedding.

Proof We proceed by induction on $\text{ntr}(A_2, \phi_2)$. If $\text{ntr}(A_2, \phi_2) = 0$ then the conclusions of the proposition hold by Proposition 6.5(fin). For $\text{ntr}(A_2, \phi_2) \geq 1$ there is a $*$ -subalgebra $B_2 \subseteq A_2$ and a partial isometry $v \in A_2$ that together with B_2 generates A_2 , such that $\text{ntr}(B_2, \phi_2|_{B_2}) < \text{ntr}(A_2, \phi_2)$, v^*v and vv^* are orthogonal minimal projections in $(A_2)_{\phi_2}$ and $(v^*v)B_2(vv^*) = \{0\}$. Indeed, just let $v \in (A_2)_{\phi_2}(\lambda^{-1})$ be a partial isometry between appropriate orthogonal minimal projections of A_2 and let $B_2 = (1 - vv^*)A_2(1 - vv^*) + \mathbf{C}v^*v$. Let $\mathcal{Q} = W^*(A_1 \cup B_2)$. Then

$$(\mathcal{Q}, \phi|_{\mathcal{Q}}) \cong (A_1, \phi_1) * (B_2, \phi_2|_{B_2}), \quad (25)$$

and by inductive hypothesis, the conclusion of the proposition applies to the free product in (25). We have $\mathcal{M} = W^*(\mathcal{Q} \cup \{v\})$. Let $p = v^*v$, $q = vv^*$ and assume without loss of generality that $0 < \lambda \stackrel{\text{def}}{=} \frac{\phi(q)}{\phi(p)} < 1$. The proof will be for three separate cases:

- (A) $\lambda \in Sd(\phi|_{\mathcal{Q}})$;
- (B) $\lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{Q}}) = \{1\}$;
- (C) $\lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{Q}}) = \lambda^{N\mathbf{Z}}$, some $N \geq 2$.

Proof in Case A There is $y \in \mathcal{Q}_\phi(\lambda^{-1})$ such that $y^*y = p$ and $yy^* = q$. Then by Lemma 5.4, it follows that

$$(p\mathcal{M}p, \phi(p)^{-1}\phi|_{p\mathcal{M}p}) \cong (p\mathcal{Q}p, \phi(p)^{-1}\phi|_{p\mathcal{Q}p}) * (L(\mathbf{Z}), \tau_{\mathbf{Z}}),$$

so by Proposition 6.5 we have that $p\mathcal{M}p$ is a type III factor, $p\mathcal{M}_\phi p \cong L(F_\infty)$ and $Sd(\phi|_{p\mathcal{M}p}) = Sd(\phi|_{p\mathcal{Q}p}) = Sd(\phi|_{\mathcal{Q}})$. Hence the conclusions of the proposition hold in Case A.

Proof in Case B There is a partial isometry $y \in \mathcal{Q}_\phi$ such that $yy^* = q$, $h \stackrel{\text{def}}{=} y^*y \leq p$. Then $p\mathcal{M}p = W^*(p\mathcal{Q}p \cup \{y^*v\})$ and $w \stackrel{\text{def}}{=} y^*v$ is a nonunitary isometry belonging to $\mathcal{M}_\phi(\lambda^{-1})$. Let

$$\begin{aligned}\mathcal{Q}_{[0,0]} &= p\mathcal{Q}p \\ \mathcal{Q}_{[-n,0]} &= W^*(w^*\mathcal{Q}_{[-n+1,0]}w \cup \mathcal{Q}_{[0,0]}), \quad (n \geq 1) \\ \mathcal{Q}_{(-\infty,0]} &= W^*(\bigcup_{n \geq 1} \mathcal{Q}_{[-n,0]}) \\ \mathcal{Q}_{[0,n]} &= W^*(w\mathcal{Q}_{[0,n-1]}w^* \cup \mathcal{Q}_{[0,0]}), \quad (n \geq 1) \\ \mathcal{Q}_{(0,\infty)} &= W^*(\bigcup_{n \geq 1} \mathcal{Q}_{[0,n]}) \\ \mathcal{Q}_{(-\infty,\infty)} &= W^*(w^*\mathcal{Q}_{(-\infty,0]}w \cup \mathcal{Q}_{[0,\infty)}).\end{aligned}$$

Then

- (i) $w^*\mathcal{Q}_{[-n+1,0]}w$ and $\mathcal{Q}_{[0,0]}$ are free in $p\mathcal{M}p \forall n \geq 1$.
- (ii) $w\mathcal{Q}_{[0,n-1]}w^*$ and $h\mathcal{Q}_{[0,0]}h$ are free in $h\mathcal{M}h \forall n \geq 1$.
- (iii) $w^*\mathcal{Q}_{(-\infty,0]}w$ and $\mathcal{Q}_{[0,\infty)}$ are free in $p\mathcal{M}p$.
- (iv) $\mathcal{Q}_{(-\infty,\infty)} = p\mathcal{M}_\phi(\Gamma)p$ where $\Gamma = Sd(\phi|_{\mathcal{Q}})$.

In order to prove (i)–(iv), we will want to keep track of occurrences of w and w^* . Hence for

$$x \in \Lambda^\circ((p\mathcal{Q}p)^\circ, \{w^n, (w^*)^n \mid n \geq 1\}), \quad (26)$$

let $l(x)$ denote the length of the word x and define $t_j(x)$ for $1 \leq j \leq l(x)$ as follows. Setting $t_0(x) = 0$ we let

$$t_j(x) = \begin{cases} t_{j-1}(x) & \text{if the } j\text{th letter of } x \text{ is from } (p\mathcal{Q}p)^\circ \\ t_{j-1}(x) + k & \text{if the } j\text{th letter of } x \text{ is } w^k \\ t_{j-1}(x) - k & \text{if the } j\text{th letter of } x \text{ is } (w^*)^k. \end{cases}$$

Let χ be the set of words x as in (26) such that any letter of x coming from $(p\mathcal{Q}p)^\circ$ and lying between w^* on the left and w on the right belongs to $(h\mathcal{Q}h)^\circ$.

Claim 7.3a *Let $I \subseteq \mathbf{Z}$ be an interval of the form $[-n, 0]$, $[0, n]$ (for $n \geq 1$), $(-\infty, 0]$, $[0, \infty)$ or $(-\infty, \infty)$. Then*

$$\mathcal{Q}_I = \text{span}(\{p\} \cup \chi_I) \text{ where } \chi_I = \{x \in \chi \mid t_{l(x)}(x) = 0, t_j(x) \in I \forall 1 \leq j \leq l(x)\}. \quad (27)$$

This claim is easily proved, (cf the proof of Claim 6.1a, *etcetera*).

Let us now show that $\chi_{(-\infty,\infty)} \subseteq \ker \phi$. Indeed, writing out $w = y^*v$, we see that every element $x \in \chi_{(-\infty,\infty)}$ is equal to an $x' \in \Lambda^\circ(\mathcal{Q}^\circ, \{v, v^*\})$ of which each letter coming from $\mathcal{Q}^\circ$ and

- lying between v on the left and v on the right belongs to $p\mathcal{Q}q$,
- lying between v on the left and v^* on the right belongs to $(p\mathcal{Q}p)^\circ$,
- lying between v^* on the left and v on the right belongs to $(q\mathcal{Q}q)^\circ$,
- lying between v^* on the left and v^* on the right belongs to $q\mathcal{Q}p$.

Now using the conditional expectation of $\mathcal{Q}$ onto B_2 , using that p and q are minimal projections in B_2 and that $pB_2q = \{0\}$, we see that every element of $p\mathcal{Q}q \cup (p\mathcal{Q}p)^\circ \cup (q\mathcal{Q}q)^\circ \cup q\mathcal{Q}p$ is the s.o.-limit of a bounded net in

$\Lambda^\circ(A_1^\circ, B_2^\circ) \setminus B_2^\circ$. Thus, regrouping x' by sticking v and v^* to B_2° whenever possible, using that $p, q \in (A_2)_{\phi_2}$, we see that x' equals an element of $\Lambda^\circ(A_1^\circ, A_2^\circ)$, hence by freeness $\phi(x') = 0$. Thus we have proved $\chi_{(-\infty, \infty)} \subseteq \ker \phi$.

To show (i) we need that $\phi(a) = 0$ whenever $a \in \Lambda^\circ((w^* \mathcal{Q}_{[-n+1, 0]} w)^\circ, (p \mathcal{Q} p)^\circ)$. Because w is in a spectral subspace $\mathcal{M}_\phi(\lambda^{-1})$ of $\mathcal{M}$, $(w^* \mathcal{Q}_{[-n+1, 0]} w)^\circ = w^*(h \mathcal{Q}_{[-n+1, 0]} h)^\circ w$. Using (27) and the conditional expectation of $\mathcal{Q}_{[-n+1, 0]}$ onto $\mathcal{Q}_{[0, 0]}$ (gotten from 3.4 and (i) for $n-1$), we have that every element of $(h \mathcal{Q}_{[-n+1, 0]} h)^\circ$ is the s.o.-limit of a bounded net in $\text{span}((\chi_{[-n+1, 0]} \setminus (p \mathcal{Q} p)^\circ) \cup (h \mathcal{Q} h)^\circ)$. Thus it will suffice to show $\phi(a') = 0$ whenever

$$a' \in \Lambda^\circ\left(w^*\left((\chi_{[-n+1, 0]} \setminus (p \mathcal{Q} p)^\circ) \cup (h \mathcal{Q} h)^\circ\right)w, (p \mathcal{Q} p)^\circ\right).$$

It is easily seen that $a' \in \chi_{[-n, 0]} \subseteq \chi_{(-\infty, \infty)}$, which we know is in $\ker \phi$, so (i) is proved.

To show (ii) is even easier. Substitute $\chi_{[0, n-1]}$ for $(\mathcal{Q}_{[0, n-1]})^\circ$ and get an element of $\chi_{[0, n]} \subseteq \chi_{(-\infty, \infty)}$.

The proof of (iii) is similar to that of (i). We want, however, a ϕ -preserving conditional expectation from $\mathcal{Q}_{(-\infty, 0]}$ onto $\mathcal{Q}_{[0, 0]}$, which follows by taking inductive limits of conditional expectations from $\mathcal{Q}_{[-n, 0]}$ onto $\mathcal{Q}_{[0, 0]}$, which exists by §3.

Putting together (i), (ii) and (iii), applying Proposition 6.5 repeatedly and taking inductive limits, we get that $\mathcal{Q}_{(-\infty, \infty)}$ is a type III factor, $\phi|_{\mathcal{Q}_{(-\infty, \infty)}}$ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and with $Sd(\phi|_{\mathcal{Q}_{(-\infty, \infty)}}) = Sd(\phi|_{\mathcal{Q}}) = \Gamma$. Since $w \in \mathcal{M}_\phi(\lambda^{-1})$ and $w^* \mathcal{Q}_{(-\infty, \infty)} w = \mathcal{Q}_{(-\infty, \infty)}$, using the hypothesis of Case B we see that (iv) holds and the conclusion of the proposition holds in Case B.

Proof in Case C Note that the proof of (i), (ii) and (iii) above did not depend on the assumption $\lambda^{\mathbb{Z}} \cap Sd(\phi|_{\mathcal{Q}}) = \{1\}$, but remains valid also in Case C. Using the same notation as in the above proof of (i), (ii) and (iii), we have that $\mathcal{Q}_{[0, N-1]}$ is a type III factor, $\phi|_{\mathcal{Q}_{[0, N-1]}}$ is an extremal almost periodic state with centralizer isomorphic to $L(F_\infty)$ and with $Sd(\phi|_{\mathcal{Q}_{[0, N-1]}}) = Sd(\phi|_{\mathcal{Q}})$. We have $ww^* = y^*y \in \mathcal{Q}_{[0, 0]}$ and then after $N-1$ steps of (ii) one has that $w^N(w^*)^N \in \mathcal{Q}_{[0, N-1]}$. Thus there exists a partial isometry z in the spectral subspace $(\mathcal{Q}_{[0, N-1]})_\phi(\lambda^{-N})$ such that $z^*z = p$ and $zz^* = h' \stackrel{\text{def}}{=} w^N(w^*)^N$. Then $z^*w^N \in p\mathcal{M}_\phi p$ is a unitary. We will show that in $p\mathcal{M}_\phi p$, z^*w^N is a Haar unitary and $\{z^*w^N\}$ and $\mathcal{Q}_{[0, N-1]}$ are $*$ -free. Both will be proved at once by showing $\phi(x) = 0$ whenever

$$x \in \Lambda^\circ((\mathcal{Q}_{[0, N-1]})^\circ, \{(z^*w^N)^n \mid n \geq 1\} \cup \{((w^*)^N z)^n \mid n \geq 1\}).$$

Now x is equal to

$$x' \in \Lambda^\circ(\mathcal{Q}_{[0, N-1]}, \{w^N, (w^*)^N\})$$

where each letter of x' that

lies between w^N on the left and $(w^*)^N$ on the right belongs to $(\mathcal{Q}_{[0, N-1]})^\circ$

lies between $(w^*)^N$ on the left and w^N on the right belongs to $(h' \mathcal{Q}_{[0, N-1]} h')^\circ$.

Since $h' \leq h$ and using (27) as well as the ϕ -preserving conditional expectation from $\mathcal{Q}_{[0,N-1]}$ onto $\mathcal{Q}_{[0,0]}$, it will suffice to show $\phi(x'') = 0$ whenever x'' is obtained from x' by replacing

- each letter from $(h' \mathcal{Q}_{[0,N-1]} h')^\circ$ with an arbitrary element of $(\chi_{[0,N-1]} \setminus (p \mathcal{Q} p)^\circ) \cup (h \mathcal{Q} h)^\circ$,
- each remaining letter from $(\mathcal{Q}_{[0,N-1]})^\circ$ with an arbitrary element of $\chi_{[0,N-1]}$,
- each remaining letter from $\mathcal{Q}_{[0,N-1]}$ with an arbitrary element of $\chi_{[0,N-1]} \cup \{p\}$.

Cancelling $w^*w = p$ whenever such neighbors occur, we see that x'' is equal to an element of χ , hence $\phi(x'') = 0$. Let $\mathcal{P} = W^*(\mathcal{Q}_{[0,N-1]} \cup \{z^*w^N\})$. Then using Proposition 6.5, we see that $\mathcal{P}$ is a type III factor, $\phi|_{\mathcal{P}}$ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and with $Sd(\phi|_{\mathcal{P}}) = Sd(\phi|_{\mathcal{Q}})$. Moreover, $\mathcal{M} = W^*(\mathcal{P} \cup \{w\})$ and $w^N \in \mathcal{P}$. From this and the assumption of Case C, one easily proves $\mathcal{M}_\phi = \mathcal{P}_\phi$ and that the conclusions of the proposition hold in Case C. □

Definition 7.4 We define the notation we shall use for faithful states on finite dimensional algebras (and other algebras). Consider a pair $(\mathcal{M}, \phi)$ where $\mathcal{M}$ is a von Neumann algebra of the form

$$\mathcal{M} = D \quad \text{or} \quad \mathcal{M} = \mathcal{Q} \oplus D,$$

with D finite dimensional and where ϕ is a normal, faithful state on $\mathcal{M}$. To describe D and the functional $\phi|_D$, we will use the following notation. For $\mathcal{M} = \mathcal{Q} \oplus D$, we write

$$(D, \phi|_D) = \bigoplus_{j=1}^K \bigoplus_{\alpha_{j,1}, \dots, \alpha_{j,n_j}}^{p_{j,1}, \dots, p_{j,n_j}} M_{n_j}(\mathbf{C})$$

which indicates $D = \bigoplus_{j=1}^K M_{n_j}(\mathbf{C})$ and

$$\phi|_D = \bigoplus_{j=1}^K \phi_j \quad \text{with} \quad \phi_j = \text{Tr}_{n_j} \left(\cdot \begin{pmatrix} \alpha_{j,1} & & 0 \\ & \ddots & \\ 0 & & \alpha_{j,n_j} \end{pmatrix} \right),$$

where Tr_{n_j} is the unnormalized trace on $M_{n_j}(\mathbf{C})$. By $p_{j,i}$ we denote the projection $\text{diag}(0, \dots, 0, 1, 0, \dots, 0)$ with 1 in the i th place. We must have that $\alpha_{j,i} > 0 \forall j, i$ and, if $\mathcal{M} = D$, that $\sum_{j=1}^K \sum_{i=1}^{n_j} \alpha_{j,i} = 1$. If $\mathcal{M} = \mathcal{Q} \oplus D$ then we may write

$$(\mathcal{M}, \phi) = \bigoplus_{\alpha_0}^{p_0} \bigoplus_{j=1}^K \bigoplus_{\alpha_{j,1}, \dots, \alpha_{j,n_j}}^{p_{j,1}, \dots, p_{j,n_j}} M_{n_j}(\mathbf{C})$$

where $p_0 = 1_{\mathcal{Q}} \oplus 0$ and $\phi(p_0) = \alpha_0$. Since ϕ is a state, we have $\alpha_0 + \sum_{j=1}^K \sum_{i=1}^{n_j} \alpha_{j,i} = 1$.

The following proposition contains Theorem 1 of the introduction.

Proposition 7.5 *Let*

$$(\mathcal{M}, \phi) = (A_1, \phi_1) * (A_2, \phi_2)$$

where for each $\iota = 1, 2$, $A_\iota \neq \mathbf{C}$ is a separable von Neumann algebra of the form D_ι , $\mathcal{Q}_\iota$ or $\mathcal{Q}_\iota \oplus D_\iota$, where D_ι is finite dimensional, $\mathcal{Q}_\iota$ is a type III factor, ϕ_ι is a normal, faithful state on A_ι and, whenever $A_\iota = \mathcal{Q}_\iota$ or $\mathcal{Q}_\iota \oplus D_\iota$, $\phi_\iota|_{\mathcal{Q}_\iota}$ is extremal almost periodic, with centralizer isomorphic to $L(F_\infty)$ or the hyperfinite II_1 factor R .

Then

$$\mathcal{M} = \mathcal{M}_0 \quad \text{or} \quad \mathcal{M} = \mathcal{M}_0 \oplus D, \quad (28)$$

where D is finite dimensional. If A_1 and A_2 both have linear dimension two, then $\mathcal{M}_0 = L(\mathbf{Z}) \otimes M_2(\mathbf{C})$. If ϕ_1 and ϕ_2 are traces, (which implies $A_1 = D_1$ and $A_2 = D_2$), and at least one of A_1 and A_2 has dimension ≥ 3 , then $\mathcal{M}_0 = L(F_t)$, some $1 < t < \infty$. If at least one of ϕ_1 and ϕ_2 is not a trace, then $\mathcal{M}_0$ is a type III factor, $\phi|_{\mathcal{M}_0}$ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and with $\text{Sd}(\phi|_{\mathcal{M}_0})$ equal to the subgroup of $\mathbf{R}_+^*$ generated by

$$\text{point spectrum}(\Delta_{\phi_1}) \cup \text{point spectrum}(\Delta_{\phi_2}).$$

It remains to decide whether D is present in (28), and in that case to describe it. If $A_1 = \mathcal{Q}_1$ or $A_2 = \mathcal{Q}_2$, then D is not present, i.e. $\mathcal{M} = \mathcal{M}_0$. Otherwise, with notation as in Definition 7.4, suppose

$$\begin{aligned} D_1 &= \bigoplus_{j=1}^{K_1} M_{n_j}(\mathbf{C}) \\ &\quad \alpha_{j,1}, \dots, \alpha_{j,n_j} \\ D_2 &= \bigoplus_{j=1}^{K_2} M_{m_j}(\mathbf{C}) \\ &\quad \beta_{j,1}, \dots, \beta_{j,m_j} \end{aligned} \quad (29)$$

If $n_j \geq 2 \forall 1 \leq j \leq K_1$ and $m_j \geq 2 \forall 1 \leq j \leq K_2$, then D is not present, i.e. $\mathcal{M} = \mathcal{M}_0$. If some $n_j = 1$ or some $m_j = 1$, we may suppose without loss of generality that $n_1 = 1$ and that $p_{1,1}$ is the largest (with respect to ϕ) of the projections in D_1 and D_2 that are both minimal and central, i.e. we suppose that

$$\alpha_{1,1} = \sup(\{\alpha_{j,1} \mid 1 \leq j \leq K_1, n_j = 1\} \cup \{\beta_{j,1} \mid 1 \leq j \leq K_2, m_j = 1\}).$$

Let

$$J = \left\{ 1 \leq j \leq K_2 \mid \sum_{i=1}^{m_j} \frac{1}{\beta_{j,i}} < \frac{1}{1 - \alpha_{1,1}} \right\}.$$

If J is empty, then $\mathcal{M} = \mathcal{M}_0$. If J is nonempty then

$$\mathcal{M} = \mathcal{M}_0 \oplus \bigoplus_{j \in J} M_{m_j}(\mathbf{C}) \quad (30)$$

where

$$\begin{aligned} \gamma_{j,i} &= \beta_{j,i} \left(1 - (1 - \alpha_{1,1}) \sum_{t=1}^{m_j} \frac{1}{\beta_{j,t}} \right) \\ r_{j,i} &= \bigwedge_{t=1}^{m_j} v_{i,t}^{(j)} (p_{1,1} \wedge q_{j,t}) v_{t,i}^{(j)}, \end{aligned}$$

where $(v_{s,t}^{(j)})_{1 \leq s, t \leq m_j}$ is a system of matrix units for the j th summand, $\mathcal{M}_{m_j}(\mathbf{C})$, of D_2 such that $v_{t,t}^{(j)} = q_{j,t}$.

Remark 7.6 We have that $(v_{s,1}^{(j)} r_{j,1} v_{1,t}^{(j)})_{1 \leq s, t \leq m_j}$ is a system of matrix units for summand j of (30), which is equal to the meet (see Dykema [1995(b)]), $\{p_{1,1}\} \odot \{v_{s,t}^{(j)} \mid 1 \leq s, t \leq m_j\}$, of matrix units of the summand of D_1 and the summand of D_2 that were mated to produce the j th summand of (30).

Proof of Theorem 7.5 The cases

$$A_1 = \mathcal{Q}_1 \text{ and } A_2 = \mathcal{Q}_2$$

$$A_1 = \mathcal{Q}_1 \text{ and } A_2 = D_2$$

$$A_1 = \mathcal{Q}_1 \text{ and } A_2 = \mathcal{Q}_2 \oplus D_2$$

can be proved by variously applying Propositions 6.5, 7.3 and 5.1. In the remaining cases, *i.e.*

$$A_1 = \mathcal{Q}_1 \oplus D_1 \text{ or } A_1 = D_1 \quad \text{and} \quad A_2 = \mathcal{Q}_2 \oplus D_2 \text{ or } A_2 = D_2, \quad (31)$$

the proposition will be proved by induction on $\text{ntr}(D_1, \phi_1|_{D_1}) + \text{ntr}(D_2, \phi_2|_{D_2})$. If the proposition has been proved in the case of A_1 and A_2 both finite dimensional, for a given value of $\text{ntr}(D_1, \phi_1|_{D_1}) + \text{ntr}(D_2, \phi_2|_{D_2})$, then it can be shown to hold also in the remaining cases of (31) by using Propositions 5.1, 6.5 and 7.3.

Hence we set about proving the case when $A_1 = D_1$ and $A_2 = D_2$ by induction. If $\text{ntr}(A_1, \phi_1) + \text{ntr}(A_2, \phi_2) = 0$ then ϕ_1 and ϕ_2 are traces and the proposition was in this case proved in Dykema [1993(b)]. Otherwise, taking $\text{ntr}(A_2, \phi_2) \geq 1$ and arguing as in the beginning of the proof of (7.3), there are $0 < \lambda < 1$, $v \in (A_2)_{\phi_2}(\lambda^{-1})$ such that $h = v^*v$ and $k = vv^*$ are orthogonal minimal projections of A_2 , and a $*$ -subalgebra $B_2 \subseteq A_2$ which together with v generates A_2 and such that h is central in B_2 and $\text{ntr}(B_2, \phi_2|_{B_2}) < \text{ntr}(A_2, \phi_2)$. Let $\mathcal{N} = W^*(A_1 \cup B_2)$. By inductive hypothesis,

$$\mathcal{N} = \mathcal{N}_0^{r_0} \quad \text{or} \quad \mathcal{N} = \mathcal{N}_0^{r_0} \oplus \bigoplus_{j \in J} M_{l_j}(\mathbf{C})^{r'_{j,1}, \dots, r'_{j,l_j}}$$

with $\mathcal{N}_0$ nonatomic and either $\phi|_{\mathcal{N}_0}$ a trace or $\phi|_{\mathcal{N}_0}$ extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and with $Sd(\phi|_{\mathcal{N}_0})$ equal to the subgroup of $\mathbf{R}_+^*$ generated by point spectrum(Δ_{ϕ_1}) $\cup$ point spectrum($\Delta_{\phi_2|_{B_2}}$). Clearly $(h+k)\mathcal{M}(h+k) = W^*((h+k)\mathcal{N}(h+k) \cup \{v\})$. Thus $\mathcal{C}_{\mathcal{M}}(h+k) = \mathcal{C}_{\mathcal{N}}(h+k)$ and once we have determined $(h+k)\mathcal{M}(h+k)$, we will be able to find $\mathcal{M}$.

One of the following four cases must hold:

- (I) $k \leq r_0$ and $h \leq r_0$;
- (II) $k \leq r_0$ and $h \not\leq r_0$;
- (III) $k \not\leq r_0$ and $h \leq r_0$;
- (IV) $k \not\leq r_0$ and $h \not\leq r_0$.

By the fact that h is central in B_2 and by inductive hypothesis, we see that case III cannot occur. We will show in cases I and II that $(h+k)\mathcal{M}(h+k)$ is a type III factor, meaning that h and k contribute only to the type III part of $\mathcal{M}$, whereas in case IV, if h and k are large enough then they can contribute to a type I direct summand of $\mathcal{M}$.

Case I is treated in three separate subcases:

- (Ia) $\mathcal{N}_0$ is a type III factor and $\lambda \in Sd(\phi|_{\mathcal{N}_0})$;
- (Ib) $\phi|_{\mathcal{N}_0}$ is a trace or $\mathcal{N}_0$ is a type III factor and $\lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{N}_0}) = \{1\}$;
- (Ic) $\mathcal{N}_0$ is a type III factor and $\lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{N}_0}) = \lambda^{N\mathbf{Z}}$, some $N \geq 2$.

The proposition in each of these cases is proved like the corresponding case in the proof of Proposition 7.3.

In Case II,

$$(h+k)\mathcal{N}(h+k) = (h+k)\mathcal{N}_0(h+k) \oplus \bigoplus_{j \in J'}^{r'_{j,1}, \dots, r'_{j,n_j}} M_{n_j}(\mathbf{C}),$$

with $\emptyset \neq J' \subseteq \{1, \dots, K_1\}$ and $r'_{j,i} \leq h \forall (j, i)$. We will now show that $\phi(r_0 h) \geq \phi(k)$. If h is the largest central and minimal projection in A_1 and B_2 , then

$$\begin{aligned} \phi(r'_{j,i}) &= \alpha_{j,i} \left(1 - (1 - \phi(h)) \sum_{t=1}^{n_j} \frac{1}{\alpha_{j,t}} \right) \\ &\leq \alpha_{j,i} \left(1 - (1 - \phi(h)) \frac{1}{\alpha_{j,i}} \right) \\ &= \phi(h) + \alpha_{j,i} - 1. \end{aligned}$$

Let $N = \sum_{j \in J'} n_j$. If $N = 1$ then $J' = \{j\}$ and $\phi(r_0 h) = \phi(h) - \phi(r'_{j,1}) \geq 1 - \alpha_{j,1} \geq 1 - \phi(h) \geq \phi(k)$. If $N \geq 2$ then

$$\begin{aligned} \phi(r_0 h) &= \phi(h) - \sum_{j \in J', 1 \leq i \leq n_j} \phi(r'_{j,i}) \\ &\geq \phi(h) - \sum_{j \in J', 1 \leq i \leq n_j} (\phi(h) + \alpha_{j,i} - 1) \\ &\geq (N-1)(1 - \phi(h)) \geq 1 - \phi(h) \geq \phi(k). \end{aligned}$$

If h is not the largest central and minimal projection in A_1 and B_2 , then, by inductive hypothesis and the assumption on the form of $\mathcal{N}_0$, the largest central and minimal projection in A_1 and B_2 must be $p = p_{j,1} \in A_1$. Hence $J' = \{j\}$, $n_j = 1$, $r'_{j,1} = p \wedge h$, $\phi(r'_{j,1}) = \phi(p) + \phi(h) - 1$ and $\phi(p) + \phi(k) < 1$. Thus $\phi(r_0 h) = 1 - \phi(p) > \phi(k)$.

Consider the three subcases:

(IIa) $\lambda \in Sd(\phi|_{\mathcal{N}});$

(IIb) $\lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{N}}) = \{1\};$

(IIc) $\lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{N}}) = \lambda^{N\mathbf{Z}}, \text{ some } N \geq 2.$

In Case IIa, there is $y \in \mathcal{N}_\phi(\lambda^{-1})$ such that $h' \stackrel{\text{def}}{=} r_0 h = y^* y$ and $k' \stackrel{\text{def}}{=} y y^* \leq k$, thus

$$h\mathcal{M}h = W^*(v^* \mathcal{N} v \cup (h - h') \mathcal{N} (h - h') \cup \{y^* v\}).$$

Let

$$\mathcal{P} = W^*(v^* \mathcal{N} v \cup (h - h') \mathcal{N} (h - h')).$$

One easily shows that $v^* \mathcal{N} v$ and $\mathbf{C}h' + (h - h') \mathcal{N} (h - h')$ are free in $(h\mathcal{M}h, \phi(h)^{-1} \phi|_{h\mathcal{M}h})$, so by Proposition 7.3 $\mathcal{P}$ is a type III factor and $\phi|_{\mathcal{P}}$ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and with $Sd(\phi|_{\mathcal{P}}) = Sd(\phi|_{\mathcal{N}})$. Now $v^* y y^* v = v^* k' v$ and $y^* v v^* y = h'$, and there is a partial isometry $z \in \mathcal{P}_\phi$ such that $z^* z = h'$ and $z z^* = v^* k' v$. Let $w = y^* v z$. Then $h' \mathcal{M} h' = W^*(h' \mathcal{P} h' \cup \{w\})$ and we claim that w is a Haar unitary and that $\{w\}$ and $h' \mathcal{P} h'$ are $*$ -free in $h' \mathcal{M} h'$ with respect to the restriction of ϕ . Clearly w is unitary in $h' \mathcal{M} h'$. It will suffice to show that $\phi(x) = 0$ whenever

$$x \in \Lambda^\circ((h' \mathcal{P} h')^\circ, \{w^n \mid n \in \mathbf{N}\} \cup \{(w^*)^n \mid n \in \mathbf{N}\}).$$

Writing $w = y^*vz$ and $w^* = z^*v^*y$ and then sticking z and z^* to letters from $(h'\mathcal{P}h')^\circ$ whenever possible, we see that

$$x = x' \in \Lambda^\circ(\{y^*v, v^*y\}, (h'\mathcal{P}h')^\circ \cup h'\mathcal{P}(v^*k'v) \cup (v^*k'v)\mathcal{P}h' \cup ((v^*k'v)\mathcal{P}(v^*k'v))^\circ),$$

where each letter of x' that is y^*v

is either the first letter of x' or has to the left a letter from $(h'\mathcal{P}h')^\circ$ or $(v^*k'v)\mathcal{P}h'$

and

has to the right a letter from $(v^*k'v)\mathcal{P}h'$ or $((v^*k'v)\mathcal{P}(v^*k'v))^\circ$,

and where each letter of x' that is v^*y

has to the left a letter from $h'\mathcal{P}(v^*k'v)$ or $((v^*k'v)\mathcal{P}(v^*k'v))^\circ$

and

is either the last letter of x' or has to the right a letter from $(h'\mathcal{P}h')^\circ$ or $h'\mathcal{P}(v^*k'v)$.

Each element of

$\mathcal{P}$ is the s.o.-limit of a bounded sequence in $\text{span}(\{h\} \cup \Omega)$

$(h'\mathcal{P}h')^\circ$ is the s.o.-limit of a bounded sequence in $\text{span}(h'\Omega_{l,r}h')$

$h'\mathcal{P}(v^*k'v)$ is the s.o.-limit of a bounded sequence in $\text{span}(\{h'\} \cup h'\Omega_l)$

$(v^*k'v)\mathcal{P}h'$ is the s.o.-limit of a bounded sequence in $\text{span}(\{h'\} \cup \Omega_r h')$

$((v^*k'v)\mathcal{P}(v^*k'v))^\circ$ is the s.o.-limit of a bounded sequence in $\text{span}((\Omega \setminus (v^*\mathcal{N}v)^\circ) \cup v^*(k'\mathcal{N}k')^\circ v)$,

where

$$\Omega = \Lambda^\circ((v^*\mathcal{N}v)^\circ, (\mathbf{C}h' + (h - h')\mathcal{N}(h - h'))^\circ)$$

Ω_l is the set of words belonging to Ω whose first letter belongs to $(v^*\mathcal{N}v)^\circ$

Ω_r is the set of words belonging to Ω whose last letter belongs to $(v^*\mathcal{N}v)^\circ$

$$\Omega_{l,r} = \Omega_l \cap \Omega_r.$$

Hence, by substituting these into x' , we see that in order to show that $\phi(x') = 0$ it will suffice to show that $\phi(x'') = 0$ whenever

$$x'' \in \Lambda^\circ(\{y^*v, v^*y\}, \{h\} \cup \Omega),$$

and when each letter of x'' that

has y^*v on the left and y^*v on the right belongs to $\{h\} \cup \Omega_r$

has y^*v on the left and v^*y on the right belongs to $(\Omega \setminus (v^*\mathcal{N}v)^\circ) \cup v^*(k'\mathcal{N}k')^\circ v$

has v^*y on the left and y^*v on the right belongs to $\Omega_{l,r}$

has v^*y on the left and v^*y on the right belongs to $\{h\} \cup \Omega_l$.

Writing out all letters from Ω , using that $(v^*\mathcal{N}v)^\circ = v^*(k'\mathcal{N}k')^\circ v$ and that $y \in k'\mathcal{N}h'$, then cancelling v^*v to h and vv^* to k whenever possible, we see that

$$x'' = x^{(3)} \in \Lambda^\circ(\{v, v^*\}, (h\mathcal{N}h)^\circ \cup h\mathcal{N}k \cup k\mathcal{N}h \cup (k\mathcal{N}k)^\circ).$$

Each element of $\mathcal{N}$ is the s.o.-limit of a bounded sequence in $\text{span}(\{1\} \cup \Psi)$, where $\Psi = \Lambda^\circ(A_1^\circ, B_2^\circ)$. Using the ϕ -preserving conditional expectation from $\mathcal{N}$ onto B_2 , and the facts that $k\mathcal{N}h \subseteq \mathcal{N}^\circ$, that h and k are minimal projections in B_2 and that $kB_2h = \{0\}$, we have that each element of $(k\mathcal{N}k)^\circ \cup k\mathcal{N}h \cup h\mathcal{N}k \cup (h\mathcal{N}h)^\circ$ is the s.o.-limit of a bounded sequence in $\text{span}(\Psi \setminus B_2^\circ)$. Thus it suffices to show $\phi(x^{(4)}) = 0$ for every

$$x^{(4)} \in \Lambda^\circ(\Psi \setminus B_2^\circ, \{v, v^*\}).$$

Since $B_2vB_2 \subseteq A_2^\circ$, we have that

$$x^{(4)} = x^{(5)} \in \Lambda^\circ(A_1^\circ, A_2^\circ),$$

so by freeness $\phi(x^{(5)}) = 0$. This completes the proof that w is a Haar unitary and that $\{w\}$ and $h'\mathcal{P}h'$ are free in $h'\mathcal{M}h'$.

Thus, applying Proposition 6.5 and using that the central support of h' in $\mathcal{M}$ is 1, we see that $\mathcal{M}$ is a type III factor, that ϕ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and $Sd(\phi) = Sd(\phi|_{\mathcal{N}})$. This completes the proof in Case IIa.

In Case IIb, there is a partial isometry, $y \in \mathcal{N}_\phi$, such that $y^*y \leq h' \stackrel{\text{def}}{=} r_0h$ and $yy^* = k$. Let $w = y^*v$. Then

$$h\mathcal{M}h = W^*(h\mathcal{N}h \cup \{w\}).$$

Let

$$\begin{aligned} \mathcal{N}_{[0,0]} &= h\mathcal{N}h \\ \mathcal{N}_{[-n,0]} &= W^*(w^*\mathcal{N}_{[-n+1,0]}w \cup \mathcal{N}_{[0,0]}) \quad (n \geq 1) \\ \mathcal{N}_{(-\infty,0]} &= W^*\left(\bigcup_{n \geq 0} \mathcal{N}_{[-n,0]}\right) \\ \mathcal{N}_{[0,n]} &= W^*(\mathcal{N}_{[0,0]} \cup w\mathcal{N}_{[0,n-1]}w^*) \quad (n \geq 1) \\ \mathcal{N}_{[0,\infty)} &= W^*\left(\bigcup_{n \geq 0} \mathcal{N}_{[0,n]}\right) \\ \mathcal{N}_{(-\infty,\infty)} &= W^*(w^*\mathcal{N}_{(-\infty,0]}w \cup \mathcal{N}_{[0,\infty)}). \end{aligned}$$

Just as in the proof of Proposition 7.3, one can show that

- (i) $w^*\mathcal{N}_{[-n+1,0]}w$ and $\mathcal{N}_{[0,0]}$ are free in $h\mathcal{M}h \ \forall n \geq 1$.
- (ii) $w\mathcal{N}_{[0,n-1]}w^*$ and $h\mathcal{N}_{[0,0]}h$ are free in $h\mathcal{M}h \ \forall n \geq 1$.
- (iii) $w^*\mathcal{N}_{(-\infty,0]}w$ and $\mathcal{N}_{[0,\infty)}$ are free in $h\mathcal{M}h$.
- (iv) $\mathcal{N}_{(-\infty,\infty)} = h\mathcal{M}_\phi(\Gamma)h$ where $\Gamma = Sd(\phi|_{\mathcal{N}})$.

Now repeatedly using previously (by inductive hypothesis) proven cases of this theorem, together with the notion of standard embeddings, we prove that $\mathcal{M}$ is a type III factor, that ϕ is extremal almost periodic with centralizer

isomorphic to $L(F_\infty)$ and with $Sd(\phi)$ equal to the subgroup of $\mathbf{R}_+^*$ generated by $Sd(\phi|_{\mathcal{N}})$ and λ . This finishes the proof in Case IIb.

In Case IIc, note that the facts (i), (ii) and (iii) that we proved above are more generally valid. Hence we prove that $\mathcal{N}_{[-N+1,0]}$ is a type III factor, that $\phi|_{\mathcal{N}_{[-N+1,0]}}$ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and with $Sd(\phi|_{\mathcal{N}_{[-N+1,0]}}) = Sd(\phi|_{\mathcal{N}})$. There is $z \in \mathcal{N}_\phi(\lambda^{-N})$ such that $z^*z = h$ and $zz^* = w^N(w^*)^N$. Let $u = z^*w^N$. Then as in the proof of Case C of Proposition 7.3, we see that $h\mathcal{M}(Sd(\phi|_{\mathcal{N}}))h = W^*(\mathcal{N}_{[-N+1,0]} \cup \{u\})$, that u is a Haar unitary in $h\mathcal{M}h$ and that $\{u\}$ and $\mathcal{N}_{[-N+1,0]}$ are free in $h\mathcal{M}h$ with respect to the restriction of ϕ . Hence we prove that $\mathcal{M}$ is a type III factor, that ϕ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and that $Sd(\phi)$ is the subgroup of $\mathbf{R}_+^*$ generated by $Sd(\phi|_{\mathcal{N}})$ and λ . This finishes the proof in Case IIc.

In Case IV, by inductive hypothesis there must be a central and minimal projection $p \in A_1$ such that $\phi(p) > \phi(p')$ whenever p' is a central and minimal projection in either A_1 or B_2 . Hence,

$$(h+k)\mathcal{N}(h+k) = (h+k)\mathcal{N}_0(h+k) \oplus \overset{(h+k)r_0}{\underset{\gamma_1/\phi(h+k)}{\mathbf{C}^{r_1}}} \oplus \overset{r_2}{\underset{\gamma_2/\phi(h+k)}{\mathbf{C}}}$$

with $r_1 \leq h$ and $r_2 \leq k$. By the assumption that h is central in B_2 we have

$$\begin{aligned} r_1 &= p \wedge h, & \gamma_1 &= \phi(r_1) = \phi(p) + \phi(h) - 1 \\ r_2 &\leq p \wedge k, & \gamma_2 &= \phi(r_2) \leq \phi(p) + \phi(k) - 1. \end{aligned}$$

Hence $\phi(hr_0) = 1 - \phi(p) \leq \phi(kr_0)$ and there exists $y \in \mathcal{N}_\phi$ such that $y^*y = hr_0$ and $yy^* \leq kr_0$. Thus $(h+k)\mathcal{N}(h+k) = W^*(k\mathcal{N}k \cup \{y, r_1\})$ and using $h = y^*y + r_1$ we see that

$$k\mathcal{M}k = W^*(k\mathcal{N}k \cup \{vy^*, vr_1v^*\}).$$

Let $\mathcal{P} = W^*(k\mathcal{N}k \cup \{vr_1v^*\})$. We claim that $k\mathcal{N}k$ and $\{vr_1v^*\}$ are free in $k\mathcal{M}k$. Indeed, one can show the stronger fact that $k\mathcal{N}k$ and $v\mathcal{N}v^*$ are free using that $\text{span}(\{1\} \cup \Lambda^\circ(A_1^\circ, B_2^\circ))$ is dense in $\mathcal{N}$ and that k and h are minimal projections in B_2 . Thus we have that

$$(\mathcal{P}, \phi(k)^{-1}\phi|_{\mathcal{P}}) \cong (k\mathcal{N}_0k \oplus \overset{r_2}{\underset{\gamma_2/\phi(k)}{\mathbf{C}}}) * \left(\overset{k-vr_1v^*}{\underset{1-\gamma_1/\phi(h)}{\mathbf{C}}} \oplus \overset{vr_1v^*}{\underset{\gamma_1/\phi(h)}{\mathbf{C}}} \right).$$

We see that $\frac{\gamma_2}{\phi(k)} < \frac{\gamma_1}{\phi(h)}$ and hence

$$\mathcal{P} = \begin{cases} \mathcal{P}_0 \oplus \overset{r_2 \wedge vr_1v^*}{\underset{(\gamma_2/\phi(k)) + (\gamma_1/\phi(h)) - 1}{\mathbf{C}}} & \text{if } \frac{\gamma_2}{\phi(k)} + \frac{\gamma_1}{\phi(h)} > 1 \\ \mathcal{P}_0 & \text{otherwise,} \end{cases}$$

where $\mathcal{P}_0$ is nonatomic and where either $\phi|_{\mathcal{P}_0}$ is a trace or $\mathcal{P}_0$ is a type III factor, $\phi|_{\mathcal{P}_0}$ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and $Sd(\phi|_{\mathcal{P}_0}) = Sd(\phi|_{\mathcal{N}_0})$.

We have $k\mathcal{M}k = W^*(\mathcal{P} \cup \{vy^*\})$ with $vy^*vy^* = yy^* \leq kr_0$ and $vy^*vy^* = k - vr_1v^*$. Thus if $\frac{\gamma_2}{\phi(k)} + \frac{\gamma_1}{\phi(h)} > 1$ then $r_2 \wedge vr_1v^*$ is central and minimal in $k\mathcal{M}k$. Finding the remaining part of $k\mathcal{M}k$ (i.e. by adding vy^* to $\mathcal{P}_0$), is done in three subcases:

$$(IVa) \quad \lambda \in Sd(\phi|_{\mathcal{N}_0});$$

$$(IVb) \quad \lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{N}_0}) = \{1\};$$

(IVc) $\lambda^{\mathbf{Z}} \cap Sd(\phi|_{\mathcal{N}_0}) = \lambda^{N\mathbf{Z}}$ for some $N \geq 2$.

Let $\tilde{k} = k - r_2 \wedge vr_1v^*$. We will now concentrate on proving in each of these three cases that $\tilde{k}\mathcal{M}\tilde{k}$ is a type III factor, that $\phi|_{\tilde{k}\mathcal{M}\tilde{k}}$ is an extremal almost periodic state with centralizer isomorphic to $L(F_\infty)$ and with $Sd(\phi|_{\tilde{k}\mathcal{M}\tilde{k}})$ equal to the subgroup of $\mathbf{R}_+^*$ generated by $(\text{point spectrum}(\Delta_{\phi_1}) \cup \text{point spectrum}(\Delta_{\phi_2}))$.

Proof in Case IVa There is $z \in \mathcal{P}_\phi(\lambda^{-1})$ such that $z^*z = yy^*$ and $zz^* = k - vr_1v^*$. Let $w = vy^*z^*$. Then $w^*w = k - vr_1v^* = ww^*$. Let $k' = w^*w$. Then $k'\mathcal{M}k' = W^*(k'\mathcal{P}k' \cup \{w\})$.

Claim 7.5a *In $(k'\mathcal{M}k', \phi(k')^{-1}\phi|_{k'\mathcal{M}k'})$, $k'\mathcal{P}k'$ and $\{w\}$ are $*$ -free and w is a Haar unitary.*

Proof of Claim 7.5a It will suffice to show $\phi(x) = 0$ whenever

$$x \in \Lambda^\circ((k'\mathcal{P}k')^\circ, \{w^n \mid n \in \mathbf{N}\} \cup \{(w^*)^n \mid n \in \mathbf{N}\}).$$

Writing $w = (vy^*)z^*$, we see $x = x' \in \Lambda^\circ(\mathcal{P}, \{vy^*, yv^*\})$, where every letter of x that lies between

vy^* on the left and yv^* on the right belongs to $(yy^*\mathcal{P}yy^*)^\circ$

yv^* on the left and vy^* on the right belongs to $(k'\mathcal{P}k')^\circ$.

Now every element of

$$\begin{aligned} \mathcal{P} & \text{ is the s.o.-limit of a bounded sequence in } \text{span}(\{k\} \cup \Theta), \\ (yy^*\mathcal{P}yy^*)^\circ & \text{ is the s.o.-limit of a bounded sequence in } \text{span}\Theta, \\ (k'\mathcal{P}k')^\circ & \text{ is the s.o.-limit of a bounded sequence in } \text{span}k'\Theta_{l,r}k', \end{aligned} \tag{32}$$

where

$$\Theta = \Lambda^\circ((k\mathcal{N}k)^\circ, \{k' - \frac{\phi(k')}{\phi(k)}k\})$$

and $\Theta_{l,r}$ is the set of $a \in \Theta$ whose first and last letters come from $(k\mathcal{N}k)^\circ$.

Replacing elements of $\mathcal{P}$ with elements of $\{k\} \cup \Theta$ and $k'\Theta_{l,r}k'$ according to (32), sticking y and y^* to elements of $(k\mathcal{N}k)^\circ$ whenever possible, and noting $y^*(k\mathcal{N}k)^\circ y \subseteq (h\mathcal{N}h)^\circ$, we see that in order to show $\phi(x') = 0$ it will suffice to show that $\phi(x'') = 0$ whenever

$$x'' \in \Lambda^\circ((k\mathcal{N}k)^\circ \cup k\mathcal{N}h \cup h\mathcal{N}k \cup (h\mathcal{N}h)^\circ, \{v, v^*\}).$$

Each element of $\mathcal{N}$ is the s.o.-limit of a bounded sequence in $\text{span}(\{1\} \cup \Psi)$, where $\Psi = \Lambda^\circ(A_1^\circ, B_2^\circ)$. Using the ϕ -preserving conditional expectation from $\mathcal{N}$ onto B_2 , and the facts that $k\mathcal{N}h \subseteq \mathcal{N}^\circ$, that h and k are minimal projections in B_2 and that $kB_2h = \{0\}$, we have that each element of $(k\mathcal{N}k)^\circ \cup k\mathcal{N}h \cup h\mathcal{N}k \cup (h\mathcal{N}h)^\circ$ is the s.o.-limit of a bounded sequence in $\text{span}(\Psi \setminus B_2^\circ)$. Thus it suffices to show $\phi(x^{(3)}) = 0$ for every

$$x^{(3)} \in \Lambda^\circ(\Psi \setminus B_2^\circ, \{v, v^*\}).$$

Since $B_2vB_2 \subseteq A_2^\circ$, we have that

$$x^{(3)} = x^{(4)} \in \Lambda^\circ(A_1^\circ, A_2^\circ),$$

so by freeness $\phi(x^{(4)}) = 0$. This completes the proof of Claim 7.5a.

Using the above claim and Proposition 7.3, then expanding out with partial isometries from $(\mathcal{P}_0)_\phi$, allows us to see the proof of the required facts about $\tilde{k}\mathcal{M}\tilde{k}$ and $\phi|_{\tilde{k}\mathcal{M}\tilde{k}}$ in Case IVa.

Proof in Case IVb Since $vy^* \in \mathcal{M}_\phi(\lambda^{-1})$, there is a partial isometry $z \in (\mathcal{P}_0)_\phi$ such that $zz^* = k - vr_1v^*$ and $z^*z \leq yy^*$. Let $w = z^*vy^*$, so that $w^*w = yy^*$ and $ww^* = z^*z \leq yy^*$. Let $k' = w^*w$, $k_1 = ww^* \leq k'$. Then $k'\mathcal{M}k' = W^*(k'\mathcal{P}k' \cup \{w\})$. Let

$$\begin{aligned} \mathcal{P}_{[0,0]} &= k'\mathcal{P}k' \\ \mathcal{P}_{[-n,0]} &= W^*(w^*\mathcal{P}_{[-n+1,0]}w \cup \mathcal{P}_{[0,0]}), \quad (n \geq 1) \\ \mathcal{P}_{(-\infty,0]} &= W^*(\bigcup_{n \geq 1} \mathcal{P}_{[-n,0]}) \\ \mathcal{P}_{[0,n]} &= W^*(w\mathcal{P}_{[0,n-1]}w^* \cup \mathcal{P}_{[0,0]}), \quad (n \geq 1) \\ \mathcal{P}_{(0,\infty]} &= W^*(\bigcup_{n \geq 1} \mathcal{P}_{[0,n]}) \\ \mathcal{P}_{(-\infty,\infty)} &= W^*(w^*\mathcal{P}_{(-\infty,0]}w \cup \mathcal{P}_{[0,\infty)}). \end{aligned}$$

Then

- (i) $w^*\mathcal{P}_{[-n+1,0]}w$ and $\mathcal{P}_{[0,0]}$ are free in $k'\mathcal{M}k' \forall n \geq 1$.
- (ii) $w\mathcal{P}_{[0,n-1]}w^*$ and $k_1\mathcal{P}_{[0,0]}k_1$ are free in $k_1\mathcal{M}k_1 \forall n \geq 1$.
- (iii) $w^*\mathcal{P}_{(-\infty,0]}w$ and $\mathcal{P}_{[0,\infty)}$ are free in $k'\mathcal{M}k'$.
- (iv) $\mathcal{P}_{(-\infty,\infty)} = k'\mathcal{M}_\phi(\Gamma)k'$ where $\Gamma = Sd(\phi|_{\mathcal{N}_0})$.

Statements (i)–(iv) are proved in the same spirit as the proof of Case IVa above and similarly to the proof of Case B of Proposition 7.3. Then Proposition 6.5 is applied repeatedly to conclude the required facts about $\tilde{k}\mathcal{M}\tilde{k}$ and $\phi|_{\tilde{k}\mathcal{M}\tilde{k}}$.

In Case IVc, the required facts about $\tilde{k}\mathcal{M}\tilde{k}$ and $\phi|_{\tilde{k}\mathcal{M}\tilde{k}}$ are proved similarly to the proof of Case C of Proposition 7.3.

After expanding out with partial isometries from the centralizer of $\phi|_{\mathcal{N}_0}$, we see that the same facts hold for $\mathcal{M}_0$ and $\phi|_{\mathcal{M}_0}$, namely, that $\mathcal{M}_0$ is a type III factor, that $\phi|_{\mathcal{M}_0}$ is extremal almost periodic with centralizer isomorphic to $L(F_\infty)$ and that $Sd(\phi|_{\mathcal{M}_0})$ is equal to the subgroup of $\mathbf{R}_+^*$ generated by (point spectrum(Δ_{ϕ_1}) $\cup$ point spectrum(Δ_{ϕ_2})). It remains to verify the size (with respect to ϕ) of the type I summand of $\mathcal{M}$ to which h and k contribute. After changing indices, we may assume that p , h , k and v , are, in the notation of (29),

$$\begin{aligned} p &= p_{1,1} \text{ with } n_1 = 1 \\ h &= q_{1,1} \text{ with } m_1 > 1 \\ k &= q_{1,2} \\ v &= v_{21}^{(1)}. \end{aligned}$$

Then

$$\begin{aligned}
\gamma_1 &= \beta_{1,1} \left(1 - (1 - \alpha_{1,1}) \frac{1}{\beta_{1,1}} \right) \\
\gamma_2 &= \beta_{1,2} \left(1 - (1 - \alpha_{1,1}) \left(\sum_{i=2}^{m_2} \frac{1}{\beta_{1,i}} \right) \right) \\
\phi(r_2 \wedge v r_1 v^*) &= \phi(k) \left(\frac{\gamma_1}{\phi(k)} + \frac{\gamma_2}{\phi(h)} - 1 \right) \\
&= \beta_{1,2} \left(1 - (1 - \alpha_{1,1}) \left(\frac{1}{\beta_{1,1}} + \sum_{i=2}^{m_2} \frac{1}{\beta_{1,i}} \right) \right),
\end{aligned}$$

which gives the correct formula for the type I summand of $\mathcal{M}$ to which h and k contribute, thus finishing the proof of the proposition in Case IV. $\square$

The following is essential for investigating free products with $B(\mathcal{H})$.

Proposition 7.7 *Let*

$$\begin{aligned}
(\mathcal{M}, \phi) &= (A_1, \phi_1) * (A_2, \phi_2) \\
&\cup \\
(\mathcal{N}, \phi|_{\mathcal{N}}) &= (B_1, \phi_1|_{B_1}) * (B_2, \phi_2|_{B_2}),
\end{aligned}$$

where ϕ_1 and ϕ_2 are faithful states, A_1 and A_2 are finite dimensional, $\mathbf{C} \neq B_1 \subsetneq A_1$, $\mathbf{C} \neq B_2 \subseteq A_2$ and at least one of $\phi_1|_{B_1}$ and $\phi_2|_{B_2}$ is not a trace. Thus by Proposition 7.5,

$$\begin{aligned}
\mathcal{M} &= \overset{r_0}{\mathcal{M}_0} \oplus D \quad \text{or} \quad \mathcal{M} = \mathcal{M}_0 \\
\mathcal{N} &= \overset{s_0}{\mathcal{N}_0} \oplus C \quad \text{or} \quad \mathcal{N} = \mathcal{N}_0
\end{aligned}$$

with $\mathcal{M}_0$ and $\mathcal{N}_0$ type III factors such that the centralizers of $\phi|_{\mathcal{M}_0}$ and $\phi|_{\mathcal{N}_0}$ are isomorphic to $L(F_\infty)$. Then $s_0 \leq r_0$ and the inclusion of centralizers,

$$(\mathcal{N}_0)_\phi \hookrightarrow s_0(\mathcal{M}_0)_\phi s_0, \tag{33}$$

is a standard embedding.

As the techniques of proof of 7.5 are similar to those of 7.3, one proves that (33) is a standard embedding like in 7.3.

Now Theorems 3 and 4 are proved by applying previously proved results and taking inductive limits.

8. Fullness of free product factors.

Murray and von Neumann [1943] defined property Gamma for factors of type II_1 in terms of the existence of nontrivial central sequences, and they showed that the von Neumann algebra of a free group, $L(F_2)$, is non-Gamma. Fullness is a property of factors that was invented by Connes [1974] and that when restricted to the II_1 situation is the property of being non-Gamma. More recently, in Theorem 11 of Barnett [1995], it was shown, for a free product of von Neumann algebras, $(\mathcal{M}, \phi) = (A, \phi_A) * (B, \phi_B)$, that $\mathcal{M}$ is full if the centralizers of ϕ_A and ϕ_B contain

enough unitaries. His result is based on the “14 ϵ Lemma” of Pukansky [1953]. From these partial results, and by analogy with the situation for free products with respect to traces, Dykema [1993(b)], one would expect that $\mathcal{M}_0$ in Theorems 1, 3 and 4 is always a full factor. Although this has not been proven in general, there are several classes of cases in which $\mathcal{M}_0$ of Theorems 1, 3 and 4 is known to be a full factor:

- (i) if $\mathcal{M}_0$ is a type III $_\lambda$ factor for $0 < \lambda < 1$ then it follows from Proposition 2.3 of Connes [1974] and the fact that the centralizer of $\phi|_{\mathcal{M}_0}$ is $L(F_\infty)$ and it thus non-Gamma, that $\mathcal{M}_0$ is full;
- (ii) in many cases, the criteria of Theorem 11 of Barnett [1995] will apply, or can be used in conjunction with Propositions 5.1 and 5.2;
- (iii) in the case of $(\mathcal{M}, \phi) = (M_2(\mathbf{C}), \phi_1) * (M_2(\mathbf{C}), \phi_2)$, for ϕ_1 and ϕ_2 arbitrary faithful states, it is proved in Dykema [SNU] that $\mathcal{M}$ is a full factor, the proof being based on Pukansky’s “14 ϵ Lemma.”

For example, using only Theorem 11 of Barnett [1995] and our Theorem 4, we can prove the existence of a full factor having arbitrary countable Sd invariant and an almost periodic state with centralizer isomorphic to $L(F_\infty)$.

Proposition 8.1 *For every countable subgroup, Γ , of $\mathbf{R}_+^*$, there is a full type III factor, $\mathcal{M}$, having extremal almost periodic state, ϕ , such that $Sd(\phi) = \Gamma$ and whose centralizer, $\mathcal{M}_\phi$, is isomorphic to $L(F_\infty)$.*

Proof For $n \geq 1$ let $0 < \lambda_n < 1$ be such that Γ is generated by $\{\lambda_n \mid n \geq 1\}$, and let

$$(\mathcal{M}, \phi) = \bigast_{n \geq 1} (M_2(\mathbf{C}), \text{Tr} \left(\begin{pmatrix} \frac{1}{1+\lambda_n} & 0 \\ 0 & \frac{\lambda_n}{1+\lambda_n} \end{pmatrix} \cdot \right)).$$

Grouping the free copies of $M_2(\mathbf{C})$ into two infinite sets and applying Theorem 4, we see that $(\mathcal{M}, \phi)$ is isomorphic to the free product of $(\mathcal{M}_1, \phi_1)$ and $(\mathcal{M}_2, \phi_2)$ with the centralizer of each ϕ_i being $L(F_\infty)$. Hence Barnett’s theorem applies to show that $\mathcal{M}$ is a full factor, while by Theorem 2, ϕ is extremal almost periodic with centralizer $L(F_\infty)$ and $Sd(\phi) = \Gamma$. $\square$

§9. Questions.

We conclude this paper with a few questions.

Question 9.1 For a given $0 < \lambda < 1$, are the type III $_\lambda$ factors that are obtainable by taking free products of various algebras isomorphic to each other? One approach to this question would be to find the trace scaling automorphism of the core factor $L(F_\infty) \otimes B(\mathcal{H})$. Rădulescu [preprint] has identified this automorphism in the case of

$$(L(\mathbf{Z}), \tau_{\mathbf{Z}}) * \left(M_2(\mathbf{C}), \text{Tr} \left(\begin{pmatrix} \frac{1}{1+\lambda} & 0 \\ 0 & \frac{\lambda}{1+\lambda} \end{pmatrix} \cdot \right) \right)$$

to be the λ -scaling automorphism that is an element of his one-parameter trace-scaling action Rădulescu [1992(b)], which he constructed via random matrices. One might hope that all type III $_\lambda$ factors constructed as free products of finite dimensional or hyperfinite algebras in this paper are isomorphic to Rădulescu’s. It may be that the approach of Dykema [1995(b)] would shed light on this.

Question 9.2 What are the continuous cores of the full type III $_1$ factors obtained in this paper as free products of finite dimensional or hyperfinite algebras? Are the continuous cores of those that have different Sd invariants

isomorphic or non-isomorphic? If they were isomorphic, then there would be uncountably many non-outer conjugate trace scaling actions of $\mathbf{R}_+^*$ on this II_∞ factor. By “continuous core” of a type III_1 factor, $\mathcal{M}$, we mean the type II_∞ factor, $\mathcal{N}$, such that $\mathcal{M} \cong \mathcal{N} \rtimes_\alpha \mathbf{R}_+^*$, where α is a trace-scaling action, (see Takesaki [1973]). The discrete core of each of these full factors is $L(F_\infty) \otimes B(\mathcal{H})$, and by Dykema [1995(b)] the continuous core is in each case isomorphic to the fixed point subalgebra of $L^\infty(\mathbf{R}_+^*, L(F_\infty) \otimes B(\mathcal{H}))$ under an certain action of the discrete group $Sd(\mathcal{M})$.

Question 9.3 Are the full type III_1 factors having the same Sd invariant that are obtainable by taking free products of various finite dimensional or hyperfinite algebras isomorphic to each other?

Question 9.4 Is there a type III_1 factor, $\mathcal{M}$, with extremal almost periodic state, ϕ , such that $\mathcal{M}_\phi$ is $L(F_\infty)$ (or any other non-Gamma II_1 factor) but $\mathcal{M}$ is not full?

References

- Arveson, W. [1974] *On groups of automorphisms of operator algebras*, J. Funct. Anal. **15**, 217-243.
- Barnett, L. [1995] *Free product von Neumann algebras of type III*, Proc. Amer. Math. Soc. **123**, 543-553.
- Connes, A. [1973] *Une classification des facteurs de type III*, Ann. scient. Éc. Norm. Sup. 4^e serie **6**, 133-252.
- Connes, A. [1974] *Almost periodic states and factors of type III_1* , J. Funct. Anal. **16**, 415-445.
- Dykema, K.J. [1993(a)] *On certain free product factors via an extended matrix model*, J. Funct. Anal. **112**, 31-60.
- Dykema, K.J. [1993(b)] *Free products of hyperfinite von Neumann algebras and free dimension*, Duke Math. J. **69**, 97-119.
- Dykema, K.J. [1994(a)] *Interpolated free group factors*, Pacific J. Math. **163**, 123-135.
- Dykema, K.J. [1994(b)] *Factoriality and Connes’ invariant $T(\mathcal{M})$ for free products of von Neumann algebras*, J. reine angew. Math. **450**, 159-180.
- Dykema, K.J. [1995(a)] *Amalgamated free products of multi-matrix algebras and a construction of subfactors of a free group factor*, Amer. J. Math. **117**, 1555-1602.
- Dykema, K.J. [1995(b)] *Crossed product decompositions of a purely infinite von Neumann algebra with faithful, almost periodic weight*, Indiana Univ. Math. J. **44**, 433-450.
- Dykema, K.J. [SNU] *Freeness and free products of von Neumann algebras*, Seoul National University GARC lecture notes, in preparation.
- Golodets, V.Ya. and Nessonov, N.I. [1987] *T-Property and nonisomorphic full factors of types II and III*, J. Funct. Anal. **70**, 80-89.
- Murray, F.J. and von Neumann, J. [1943] *Rings of operators. IV*, Ann. of Math. **44**, 716-808.
- Pedersen, G.K. and Takesaki, M. [1973] *The Radon-Nikodym theorem for von Neumann algebras*, Acta Math. **130**, 53-87.
- Popa, S. [1993] *Markov traces on universal Jones algebras and subfactors of finite index*, Invent. Math. **111**, 375-405.
- Pukansky, L. [1953] *Some examples of factors*, Publ. Math. Debrecen **3**, 135-156.
- Rădulescu, F. [1992(a)] *The fundamental group of the von Neumann algebra of a free group with infinitely many generators*, J. Amer. Math. Soc. **5**, 517-532.
- Rădulescu, F. [1992(b)] *A one parameter group of automorphisms of $L(F_\infty) \otimes B(\mathcal{H})$ scaling the trace*, C. R. Acad. Sci. Paris **t. 314 Série I**, 1027-1032.
- Rădulescu, F. [1994] *Random matrices, amalgamated free products and subfactors of the von Neumann algebra of a free group, of noninteger index*, Invent. Math. **115**, 347-389.
- Rădulescu, F. [preprint] *A type III_λ factor with core isomorphic to the von Neumann algebra of a free group, tensor $B(H)$* , preprint, UC-Berkeley, 1993.
- Reed, M. and Simon, B. [1972] *Methods of Modern Mathematical Physics, I: Functional Analysis*, Academic Press.
- Takesaki, M. [1970] *Tomita’s Theory of Modular Hilbert Algebras and its Applications*, Lecture Notes in Mathematics vol. 128, Springer-Verlag.
- Takesaki, M. [1973] *Duality for crossed products and the structure of von Neumann algebras of type III*, Acta. Math. **131**, 249-310.
- Tomiyama, J. [1957] *On the projection of norm one in W^* -algebras*, Proc. Japan Acad. **33**, 608-612.
- Voiculescu, D. [1985] *Symmetries of some reduced free product C^* -algebras*, Operator Algebras and Their Connections with Topology and Ergodic Theory, Lecture Notes in Mathematics, Volume 1132, Springer-Verlag, pp. 556-588.
- Voiculescu, D. [1990] *Circular and semicircular systems and free product factors*, Operator Algebras, Unitary Representations, Enveloping Algebras, and Invariant Theory, Progress in Mathematics, Volume 92, Birkhäuser, Boston, pp. 45-60.
- Voiculescu, D., Dykema, K.J. and Nica, A. [1992] *Free Random Variables*, CRM Monograph Series vol. 1, American Mathematical Society.